Evaluation of the Feeding Potentials of Amaranthus Stem and F. Thoningii Foliage as Supplements to Concentrate Diet for Weaner Rabbits

Yusuf Aliyu*

Federal College of Wildlife Management, Nigeria

***Corresponding author:** Yusuf Aliyu, Federal College of Wildlife Management, PMB 268, New Bussa, Niger State, Nigeria.

To Cite This Article: Yusuf Aliyu. Evaluation of the Feeding Potentials of Amaranthus Stem and F. Thoningii Foliage as Supplements to Concentrate Diet for Weaner Rabbits. Am J Biomed Sci & Res. 2019 - 1(6). AJBSR.MS.ID.000551. DOI: [10.34297/AJBSR.2019.01.000551](https://doi.org/10.34297/AJBSR.2019.01.000551)

Received: February 25, 2019 | **Published:** March 12, 2019

Abstract

This study was carried out to evaluate the feeding potentials of Amaranthus hybridus stem and Ficus thoningii foliage as supplements to concentrate diet for weaner rabbits over 10 weeks period. A total of 18 crossbred weaner rabbits with average initial weight of 630 ± 10 g were used for feeding trial in a completely randomized design experiment. The animals were divided into three groups of six animals per treatment and each treatment had triplicates of two animals per replicate. Treatment T1 was fed with concentrate only. Treatment T2 was fed with concentrate and A. hybridus stem while treatment T3 was fed with concentrate and F. thoningii foliage. Nutrient digestibility of the animals differed significantly ($P < 0.05$) between the groups. Feed intake and weight gain were higher in T2 than T1 and T3 although feed conversion was better in T1. It is therefore concluded that Amaranthus stem can be used as supplement to concentrate diet for weaner rabbits.

Introduction

In recent years interest in rabbit production in Nigeria has increased to improve supply of animal protein from fast growing animals with short gestation period and short generation interval. However, competition between man, animals and industries for conventional feed resources and high cost of compounding concentrate has become a major constraint militating against increased production. The increasing cost of conventional feed ingredients with the resultant increase in the cost of livestock products has necessitated research in the search for cheaper and readily available alternative feed ingredients for livestock feeding. Studies have shown that rabbit can thrive on a number of tropical forages supplemented with concentrate [2,3], but concentrate rations are expensive and can be a constraint to exploitation of the potential of rabbit. Therefore, a feeding strategy that will engage sparing use of concentrate and dependence on forages will be of immense benefit [1]. Such forages must be abundant, available all year round and accessible. Amaranthus hybridus is a widely known leaf vegetable, the foliage of which is utilized and usually discarded as waste. Ficus thoningii is a tree species that is widely distributed in many savanna areas of Nigeria which is also less compete for by man. Utilization of A. hybridus stem and F. thoningii foliage by animals can reduce waste and decrease

cost of production. This study was carried out to evaluate the feeding potentials of A. hybridus stem and F. thoningii foliage as supplements to concentrate diet for weaner rabbits.

Materials and Methods

This research work was conducted at the Teaching and Research Farm of Federal College of Wildlife Management, New Bussa, Niger State, Nigeria. Concentrate diet was compounded as shown in (Table 1). Eighteen crossbred weaner rabbits with average initial weight of 630 ± 10 g was used. The animals were divided into three treatment groups of six animals with each group having triplicate of two animals per replicate in a completely randomized design experiment. They were given prophylactic treatment and allowed for two weeks adjustment period followed by 10 weeks experimental period. Treatment 1 (T1) were fed with concentrate diet only while Treatment 2 (T2) were fed with concentrate and A. hybridus stem. Treatment 3 (T3) were fed with concentrate and F. thoningii foliage. Feed and water were served ad-libitum. Feed intake, weight gain and feed conversion ratio were measured. After the feeding trial, seven days digestibility study was conducted. Proximate analyses of the concentrate diet, A. hybridus stem, F. thoningii foliage as well as faecal samples were carried out according to [4]. All data collected

were subjected to statistical analysis using statistical package for social science [5] significant means were separated using Duncans Multiple Range test of the same package.

Table 1: Composition of the experimental diets.

Ingredient	Percentage (%)
Maize bran	64.61
Groundnut cake	18.39
Blood meal	3
Rice husk	10
Bone meal	3
Salt	0.5
Premix	0.5
Total	100
Calculated composition	
Crude protein (%)	18
Crude fiber (%)	11.7
Metabolizable Energy (Kcal/Kg)	2580

Results and Discussion

The results of proximate analyses of the concentrate, *A. hybridus* stem and *F. thoningii* is presented in (Table 2). Nutrient digestibility is shown in (Table 3). Dry matter (DM), crude protein (CP), ether extract (EE), Ash and Nitrogen free extract (NFE) were more digestible in animals on concentrate only than animals on concentrate and forages but crude fibre was more digestible in animals on concentrate and forages than those on concentrate only. Similarly, with the exception of ether extract, all other nutrients were more digestible in animals on *A. hybridus* than those on *F. thoningii*. This is an indication that *A. hybridus* stem is more digestible than *F. thoningii*. The reason may probably be due to some anti-nutritional factors present in *F. thoningii*. This is in time with the report of [6] who stated that a variety of anti-nutritional factors can affect digestion, reduce growth and affect body metabolism of animals. Digestibility of nutrients differ significantly ($P < 0.05$) between the treatment groups. Results of feed intake, weight gains and feed conversion ratio are presented in (Table 4). Average daily feed intake (ADFI) of animals on concentrate diet only (T1) was 43.00 g with corresponding average weight gain (AWG) of 14.00 g. ADFI of animals fed concentrate and *A. hybridus* stem (T2) was 80.50 g with concentrate and *A. hybridus* consumption of 30.20 g and 50.30 g respectively to gain 17.14g. Similarly, ADFI of animals fed concentrate and *F. thoningii* (T3) was 65.50 g with concentrate and *F. thoningii* consumption of 30.00 g and 35.50 g respectively to gain 12.86 g. ADFI and AWG were better in animals on concentrate and *A. hybridus* stem than other groups. Feed conversion ratio was lower in T1 than T2 and T3. All the values obtained for all the parameters were significantly ($P < 0.05$) different between the treatments. Although higher feed intake with corresponding higher gain were recorded in T2, feed was better converted to carcass in T3. However, T2 had higher gain and better conversion than T3. This could be due to higher CP, lower CF and succulent nature of *A. hybridus* stem compared to *F. thoningii* foliage. The observation is in line with the report of [1]. Higher CF in *F. thoningii* than in *A.*

hybridus stem may also be a factor that led to lower feed intake and lower digestibility in T3 than in T2. This observation is in line with report of [7] who reported reduced digestibility due to high levels of dietary fibre. This apart, the significantly lowest feed intake and weight gain of rabbits on T3 possibly suggests that *F. thoningii* may likely contain some anti-nutritional factors which must have affected its intake and utilization. The results of this study had indicated that rabbit can perform well when fed with concentrate diet supplemented with *A. hybridus* stem. This agrees with the conclusion of [8] who concluded that rabbits can perform well with mixed regime of forages and concentrate.

Table 2: Proximate Composition of the experimental concentrate, *A. hybridus* stem and *Ficus thoningii* foliage.

Nutrient (%)	Concentrate	<i>A. hybridus</i> stem	<i>F. thoningii</i>
Dry matter	91.7	50.48	64.32
Crude protein	17.89	10.05	9.73
Crude fibre	12.2	15.31	18.56
Ether extract	6.52	3.2	5.92
Ash	8.34	4.68	7.32
Nitrogen free extract	55.05	66.76	58.47

Table 3: Nutrient Digestibility of the experimental rabbits.

Nutrient (%)	T1	T2	T3	SEM
Dry matter	68.42 a	64.37 a	62.05 b	3.06
Crude protein	70.38 b	66.48 b	60.85 c	2.82
Crude fibre	58.1	60.29	59.92	1.01
Ether extract	63.39a	59.30 b	60.05 b	1.14
Ash	59.75 a	56.62 b	53.26 c	1.26
Nitrogen free extract	62.38	57.97	52.08 c	2.86

a, b, c, d Means in the same row with different superscripts differ significantly ($P < 0.05$)

SEM = Standard Error of Mean

Table 4: Performance characteristics of the experimental rabbits.

Parameters	T1	T2	T3
Initial weight (g)	630	620	640
Final weight (g)	1,610.00 b	1,819.80 a	1540.20 c
Total weight gain (g)	980.00 b	1,199.80 a	900.20 c
Average weight gain (g)	14.00 b	17.14 a	12.86 b
Total feed intake/day (g)	43.00 c	80.50 a	65.50 b
Average forage intake/day (g)	-	50.30 a	35.50 b
Average Concentrate intake/day (g)	43.00 a	30.20 b	30.00 b
Feed conversion ratio	3.07 c	4.70 b	5.09 a

a. b. c. d Means in the same row with different superscripts differ significantly ($P < 0.05$)

Conclusion

In conclusion, the results of this study had indicated that A. hybridus stem have potentials to support growth when fed with concentrate diet to weaner rabbits. It is therefore recommended that A. hybridus stem be used in mixed feeding regime for weaner rabbits especially during dry season when forages are scarce.

References

1. Yusuf AM, Olafadehan OA, Garba MH (2010) Evaluation of the feeding potentials of Vitellaria paradoxa, Nauclea latifolia and Terminalia macroptera foliage as supplements to concentrate feed for grower rabbits. Australian Journal of Basic and Applied Science 4(3): 429-433.
2. Odeyinka SM, Ijiyemi OC (1997) Performance of rabbits fed Leucaena leucocephala and concentrate between 9th and 25th weeks of age. Nigerian Journal of Animal Production 24(1): 51-53.
3. Shiawoya EL, Musa JAO (2003) Evaluation of the feeding potentials of mango (Mangifera indica) leaves, banana (Musaspp) leaves and Tridax (Tridax procumbens) as supplements to conventional feeds for growing rabbits. Proceeding of 8th Annual Conference of Animal Science Association of Nigeria (ASAN). pp 90-92.
4. AOAC (1990) Official Methods of Analysis. Association of Official Analytical Chemists, (15th edn), Washington, D.C, USA.
5. Statistical Package for Social Science (1980) SPSS for windows version 11.
6. Maidala AA, Doma UD, Aighigh DT (2011) Growth performance of weaner rabbits fed diets containing differently processed soyabean (Glycine max L. merril) products. Proceedings of 36th Annual conference of Nigerian Society for Animal Production (NSAP), Nigeria. pp 315-317.
7. Ezenwa, L., Bamikole, M.A. and Akinsoyinnu A.O. (1998). The feeding value of *Verano stylo* and Guinea grass hays as dry season feeds for rabbits. Proceedings of the 3rd Annual Conference of Animal Science Association of Nigeria (ASAN). September 22-24, Ikeja, Lagos, Nigeria. pp 203-205.
8. Bamgbose AM, Akinnusi FAO, Kuforiji AT, Osofowora AO, Osa AO, et al. (2003) Growth response of weaner rabbits fed concentrate/forage supplemented based diets. Proceedings of 8th Annual Conference of Animal Science Association of Nigeria (ASAN). September 15-18. Federal University of Technology, Minna, Nigeria. pp 98-99.