Using Bioinformatics to Study Homology Between Herpes Simplex Virus Type 1 Glycoprotein C Gene and Herpes Simplex Virus Type 2 Glycoprotein F Gene

Shereen F Elkholy^{1,2*}

¹Department of Biology, Prince Sattam bin Abdulaziz University, Alkharj, Kingdom of Saudi Arabia

²Department of Plant Molecular Biology, Agricultural Genetic Engineering Research Institute (AGERI), Agricultural Research Centre (ARC), Egypt.

*Corresponding author: Shereen F Elkholy, Department of Biology, College of Science and Humanities, Prince Sattam bin Abdulaziz University, Alkharj, Kingdom of Saudi Arabia

To Cite This Article: Shereen F Elkholy. Using Bioinformatics to Study Homology Between Herpes Simplex Virus Type 1 Glycoprotein C Gene and Herpes Simplex Virus Type 2 Glycoprotein F Gene. Am J Biomed Sci & Res. 2019 - 2(2). AJBSR.MS.ID.000575. DOI: [10.34297/AJBSR.2019.02.000575](https://doi.org/10.34297/AJBSR.2019.02.000575)

Received: February 19, 2019 | Published: March 29, 2019

Abstract

Bioinformatics is an interdisciplinary field mainly involving molecular genetics, computer science, mathematics, and statistics. Data intensive and large-scale biological problems are addressed from a computational point of view. The most common problems are modeling biological processes at the molecular level and making inferences from collected data. In this study we used bioinformatics as a tool to study homology between Herpes simplex virus type 1 glycoprotein C gene and Herpes simplex virus type 2 glycoprotein F gene. In the beginning we used the Gene data Bank to get the DNA sequence. Then we used a program Blast and got a percentage alignment 81%. We used protein data bank (PDB) to estimate the structure of the expressed proteins that were tertiary (three dimensional) structure that harbor both α helices and β pleated sheets. Finally, we used "PHYRE2" protein folding recognition set to visualize, analyze and manipulate the resulted 3D protein model.

Introduction

Bioinformatics is conceptualizing Biology in terms of molecules and applying "informatics techniques" (derived from disciplines such as applied math's, computer science and statistics) to understand and organize the information associated with these molecules, on a large scale. In short, bioinformatics is a management information system for molecular biology and has many practical applications. Bioinformatics is an application of techniques from computer science to problems from Biology. It is link between Computer Science and Biology. And has three aims: the first one: organizes data in a way that allows researchers to access existing information and to submit new entries as they are produced, e.g. the Protein Data bank for 3D macromolecular structures. The second aim: is to develop tools and resources that aid in the analysis of data. For example, having sequenced a protein, it is of interest to compare it with previously characterized sequences. The third aim: is to use these tools to analyze the data and interpret the results in a biologically meaningful manner.

Bioinformatics is a very broad field and it encompasses issues like mapping, sequencing, sequence comparison, gene identification, protein modeling, network databases, visualization

and ethics. It is an interdisciplinary subject that on one hand requires biological information-infrastructure building and on the other requires computation based biological research. All this depends on the large stores of experimental and derived data.

The foundation of Bioinformatics is based on the computational techniques, algorithms, artificial intelligence, database management, software engineering etc. All this leads to the development of community data resources and from this starts its applications development of the bioinformatics for analysis of the genetic data.

In this study, we used Bioinformatics to detect the homology and the percentage of alignment between Herpes simplex virus type 1 glycoprotein C gene and Herpes simplex virus type 2 glycoprotein F gene.

Herpes simplex virus

Is an enveloped double stranded DNA virus which encodes information for at least 10 glycoproteins. Three of the glycoproteins have functions which may modulate the immune response. In HSV-1, glycoprotein E (gE) and gI function as a complex to bind the Fc portion of immunoglobulin G [3,4,6,10,15,16,22]. gC binds the C3b and iC3b fragments of the third component of complement [7-

9,14,20,21,26-28]. Although gC is not strictly required for infection in cell culture [5,12,43,17,29], it plays an important role in virus attachment [11,25].

More recently, in-frame deletion and linker insertion mutants of gC-2 were used to identify regions important for C3b binding [26,27]. These studies showed that amino-terminal residues 26 through 73 of gC-2 are not involved in C3b receptor activity. In addition, three distinct regions (I, II, and III) in gC-2 are important for C3b binding. Region III has features like those of the short consensus repeat (SCR) [13,18,23, 24,28] of the human C3-C4 receptor CR1 [26].

Several studies employed a transient transfection system using the gC-2 gene cloned into the genomes of herpes simplex virus type 1 (HSV-1) and HSV-2 encode at least four different glycoproteins, gA/B, gC, gD, and gE, which are found on the surface of the infected cell and the virion [39]. Three of the four glycoproteins, gA/B, gD, and gE, have been found to be structurally similar in the two virus types, based on immunological and biochemical criteria [33-37]. For example, a recent analysis of the DNA sequences of the gD genes from HSV-1 and HSV-2 revealed that the gD proteins had an overall sequence homology of 85% (L. Lasky and D. Dowbenko, DNA, in press). Thus, it may be concluded that the primary sequences of these three glycoproteins have been relatively well conserved, since the two virus types diverged from each other. HSV-1 gC initially appeared to have no obvious homolog in HSV-2. HSV-1 gC was thought to be type specific since antibodies against this glycoprotein were found to react almost exclusively with HSV-1 gC [38]. In addition, no detectable immunological reactions could be demonstrated between HSV-1 gC and antisera made against HSV-2 [32]. A protein having the same electrophoretic mobility as HSV-1 gC has been demonstrated in HSV-2; however, it did not map collinearly with HSV-1 gC [17].

In contrast to HSV-1, HSV-2 appears to encode yet another glycoprotein, termed gF [30,31,35,40]. Although HSV-2 gF had an electrophoretic mobility which was much faster than HSV-1 gC, initial mapping studies with recombinant viruses revealed that this protein was encoded by a region of the HSV-2 genome which was approximately colinear with the gene for HSV-1 gC (35, 40). Subsequent studies with finer structural mapping revealed a much closer collinearity between the HSV-1 gC and the HSV-2 gF coding regions [41]. In addition, it has been recently demonstrated that a monoclonal antibody against HSV-2 gF cross-reacts weakly with HSV-1 gC [42] and that a polyclonal antiserum made against HSV-1 virion envelope proteins precipitates gF [40], suggesting a possible structural homology between the two glycoproteins. Thus, it appeared that a possible homolog to HSV-1 gC was the HSV-2 gF protein. The most conclusive proof of relatedness between two proteins is to demonstrate homology at the amino acid level.

In this research we used Bioinformatics to study homology between the Herpes Simplex Virus type 1 glycoprotein C gene and Herpes Simplex Virus type 2 glycoprotein F gene by using dry lab biometric analysis of biological data using software such as (BLAST, FASTA, PDB and PHYRE2).

Materials and Methods

GenBank

GenBank® is the NIH genetic sequence database, an annotated collection of all publicly available DNA sequences (Nucleic Acids Research, 2013 Jan;41(D1): D36-42). GenBank is part of the International Nucleotide Sequence Database Collaboration, which comprises the DNA Databank of Japan (DDBJ), the European Nucleotide Archive (ENA), and GenBank at NCBI. These three organizations exchange data daily.

A GenBank release occurs every two months and is available from the ftp site. The release notes for the current version of GenBank provide detailed information about the release and notifications of upcoming changes to GenBank. Release notes for previous GenBank releases are also available. GenBank growth statistics for both the traditional GenBank divisions and the WGS division are available from each release.

We used the gene bank to detect the DNA sequences of both glycoprotein C and glycoprotein F of herpes simplex virus type 1 and type 2 respectively (<https://www.ncbi.nlm.nih.gov/genbank/>).

Blast

BLAST finds regions of similarity between biological sequences. The program compares nucleotide or protein sequences to sequence databases and calculates the statistical significance.

The Basic Local Alignment Search Tool (BLAST) finds regions of local similarity between sequences. The program compares nucleotide or protein sequences to sequence databases and calculates the statistical significance of matches. BLAST can be used to infer functional and evolutionary relationships between sequences as well as help identify members of gene families.

We used the blast programmed to detect the homology between herpes simplex virus type 1 glycoprotein C and herpes simplex virus type 2 glycoprotein F and estimate the percentage similarity as well as the homology between some other strains (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>)

Protein Data Bank PDB

The protein data base is used to detect the amino acid sequences of both glycoprotein C and F and detect the homology between the 2 sequences (<https://www.ncbi.nlm.nih.gov/pdb>).

Phyre2 to analyze protein structure

Phyre2 is a suite of tools available on the web to predict and analyze protein structure, function and mutations. The focus of Phyre2 is to provide biologists with a simple and intuitive interface to state-of-the-art protein bioinformatics tools. Phyre2 replaces Phyre, in this updated protocol, we describe Phyre2, which uses advanced remote homology detection methods to build 3D models, predict ligand binding sites and analyze the effect of amino acid variants (e.g., nonsynonymous SNPs (nsSNPs) for a user's protein sequence. This protocol will guide users from submitting a protein sequence to interpreting the secondary and tertiary structure of their models, their domain composition and model quality. A range of additional available tools is described to find a protein structure in a genome, to submit large number of sequences at once and to automatically run weekly searches for proteins that are difficult

to model. A typical structure prediction will be returned between 30 min and 2 h after submission (<http://www.sbg.bio.ic.ac.uk/phyre2>).

Results

Detection of DNA sequence of glycoprotein C and F Gene (Table 1)

Table 1: Detection of DNA sequence of glycoprotein C and F Gene

<p>Herpes simplex virus type 1 glycoprotein C</p> <p>Sequence</p> <p>Accession no.</p> <p>KM279072.1</p>	<p>Atggccccggggcggggtgggccttgcctggctctgtggagcctgttggctcggggcg gggggtgtcgggggctcggaactgcctccacggggccacgatcaccgaggagcggtg acgaacgcgagcagggccccacatcggggtccccgggtcagccgagcccggaagt acccccacatcgacccaaaccccaacaatgtcacaaaaacaaacccccacggag ccggccagcccccaacaaccccaagccacctccacgccccaaagccccccacgtcc acccccgaccccaaaccaagaacaacacacccccgcaagtcgggcccggccactaaa ccccccggggcctgtgtgtgacgcggcaccattggcccgtacggctcgggggtg cagatccgatcgcggttcggaattccacccgatggagttccgctccagatatggcgt tactcatgggtccgtcccccaatcgtccggctccgacctagaggaggtcgtgacg aacatcaccgccccacgggggactcctgtgtacgacgcgccccaaacctgacggac ccccacgtctctggcgaggggccggcccggtgcccagctccgttctgtattctgtc accgggcccgtcgccacccagcggtgattatcgcgaggtgaccccgcgaccaggga atgtattacttggcctggggccggatggacagccgcacgagtacgggacgtgggtgagc gtccgatgttcgccccctgtctgacccctccagccccacgggtgatggagggtcag ccgttcaaggcgacgtgcacggcgacgcctactaccccgtaaccccggtggattggc tggttcgaggacgacgcccaggtgttaacccgggagatgacacgcagacgcacgag caccgcgaggggtaccacccgtctctaccgtgacctccgaggtgtcgccggccaggtc cccccgcgaccttcacgtccagatgacgtggcaccgcgactccgtgacattctcgga cgcaatgccacgggctggccctgtgtgctgcggccaacatcacatggaattggg gtcgggcatgtgtgtgacggcggtgctgctcccgaggcggtgacgtttgcctgttc ctgggggacgacccctcaccggcggttaagtcggcgttacggccaggagtcacgac caccggggtgtgacgtccgtccacccctgccattctgacgactacagcagtagac atctgtcggtgacgggatccggcgggattccgttctagaacaccacggcagtcac agccccaccagggaacccacgagcggcaggtgatcgaggcgatcgagtggtgggg attggaatcggggtctcgcggggggtcctgtgtaacggcgatcgtgtacgtcgtc cgacatcacagtcgcggcagcgtcatcgcggttaa</p>
<p>Herpes simplex virus type 2 glycoprotein F</p> <p>Sequence</p> <p>Accession no.</p> <p>X01456.1</p>	<p>Gtgcctgtggacgggtataaaggccagggggcagggcgcccatcactgttaggggtgta ggttgggaggtggcacaaaagcgacacccgtgtgtgtgttcgcccgggagggcggtg gtttccggcaaccctctcgtgctgcggcgccgcccacggctcttcgcccgggcccggg gctcttctgtcatggccttggacgggtggcctaaccgtgggctgtggggcctgtgtg tgggtgggtgtgtgtgtgtgtgccaatgcctccccggacgcacgataacgtgggc ccgcccggggaacgcgagcaatgcccccctcggtcccccggaaccgatccgccccga accacaccacgcccccaaccccgcaaggcgacgaaaagtaaggcctccaccgcaaa ccggccccgcccccaagaccggccccgaagacatcctcgagcccggtgcatgcaac cgccacgaccgctggcccgtacggctcggggtgcaaatccgatccggtttccaac tccaccgcacggagtcctcctcagatctggcgttatgccacggcgacggacgcgag atcggaacggcgcttagcttagaggaggtgatgtaaacgtgtcgccccccggggg caactgtgtatgacagcgcccccaacgaacggacccgacgtgatctggcgaggaggc gcccggccggcgccagcccggtgtactcggtcgtcgggcgctgggtcggcagcgg ctcatcatgaagagctgaccttggagaccagggtatgactactgggtgtggggccgg acggacggcccgctcggtacgggacctgggtgcgcttcggtgttcgcccctcgtcg ctgacctacccccacggtgtgtagggccagccgtttaaggcgacgtgacggcc gccactactaccgggcaaccgcggagttcgtgtggttcgaggaaggtcgccgggta ttcgatccggcccagatacacacgcagacgcaggagaacccgacggctttccaccgtc tccacgtgacctccggcgctcgggccaggcccccccgccaccttcacctgcag ctgactggcaccgcgactcgtgtgttctctcggcgaacggcagcgacggcatcg gtgctcccgccaaccattaccatggagttacgggcgacatcggtctgcacggcc ggctgtgtcccggagggtgacgtttgctgttctgggggacgactcctcgccggc gagaagggtggcgctcggtcccagacatcgtcgggcgccccggcaccgcccagatccg tccacctgcccgtctgtacgacgacagcaggtacatctccggtcgccgggataccg gacggaattccggtcctagacaccacggcagccaccagcccccccgccgggacccac gagcggcaggtgatccggcggtggaggggcggggatcgagtggtgtcctgtcgcg gtggttctggccgggacgcggtagtgtacctcaccacgcctcctcggtcgctatcgt cggctcggttaactcggggccgggcccggcgccgggtgtcttctttccacccttc gtccccgtaccaccacaccccccccccccccggtcccccgggcgttataagcc</p>

The above result indicates 81% homology between Herpes simplex virus type 1 glycoprotein C gene and Herpes simplex virus type 2 glycoprotein F sequence.

Score	Expect	Identities	Gaps	Strand
869 bits (470)	0	895/1106(81%)	6/1106(0%)	Plus/Plus
Query 346	gacc-gccccccctaa-ac-cccccg-gcccgTGTGGTGCAGCCGCCGACCCATTGGC	401		
Sbjct 438		497		
Query 402	CCGGTACGGCTCGCGGTGCAGATCCGATGCCGGTTTCGGAATCCACCCGCATGGAGTT	461		
Sbjct 498	CCGGTACGGCTCGCGGTGCAAAATCCGATGCCGGTTTCCCAACTCCACCCGCACGGAGTC	557		
Query 462	CCGCCTCCAGATATGGCGTTACTCCATGGGTCCGTcccccccAATCGCTCCGGCTCCCGA	521		
Sbjct 558	CCGCCTCCAGATCTGGCGTTATGCCACGGCGACGGACGCCGAGATCGGAACGGCGCCTAG	617		
Query 522	CCTAGAGGAGGTCTTGACGAACATCACCGCCCCACCCGGGGGACTCCTGGTGTACGACAG	581		
Sbjct 618	CTTAGAGGAGGTGATGTTAAACGTGTCGGCCCCGCCGGGGGCCAACTGGTGTATGACAG	677		
Query 582	CGCCCCCAACCTGACGGACCCCCACGTGCTCTGGGCGGAGGGGGCCGGCCCGGGCGCCGA	641		
Sbjct 678	CGCCCCCAACCGAACGGACCCGCACGTGATCTGGGCGGAGGGGCCCGGCCCGGGCGCCAG	737		
Query 642	CCCTCCGTTGTATTCTGTACCGGGCCGCTGCCGAC-CCAGCGGCTGATTATCGGCGAGG	700		
Sbjct 738	CCCGCGGCTGTACTCGGTCTGTGGGCCGCTG-GGTGCGCAGCGGCTCATCATCGAAGAGC	796		
Query 701	TGACGCCCGCGACCCAGGGAATGTATTACTTGGCCTGGGGCCGGATGGACAGCCCGCACG	760		
Sbjct 797	TGACCTTGGAGACCCAGGGCATGTACTACTGGGTGTGGGGCCGGACGGACCGCCCGTCCG	856		
Query 761	AGTACGGGACGTGGGTGCGCGTCCGCATGTTCCGCCCCCGCTCTCTGACCTCCAGCCCC	820		
Sbjct 857	CGTACGGGACCTGGGTGCGCGTTCGCGTGTTCGCCCTCCGTCGTGACCATCCACCCCC	916		
Query 821	ACGCGGTGATGGAGGGTCAGCCGTTCAAGGCAACGTCCACGGCCGCCGCTACTACCCGC	880		
Sbjct 917	ACGCGGTGCTGGAGGGCCAGCCGTTTAAGGCGACGTGCACGGCCGCCACCTACTACCCGG	976		
Query 881	GTAACCCCTCGAGTTTGTCTGGTTCGAGGACGACCGCCAGGTGTTTAAACCGGGCCAGA	940		
Sbjct 977	GCAACCGCGCGAGTTCTGTCTGGTTCGAGGACGGTCGCCGGGTATTGATCCGCGCCAGA	1036		
Query 941	TCGACACGCAGACGCACGACACCCCGACGGGTTCAACACAGTCTCTACCGTGACCTCCG	1000		
Sbjct 1037	TACACACGCAGACGCAGGAGAACCCCGACGGCTTTTCCACCGTCTCCACCGTGACCTCCG	1096		
Query 1001	AGGCTGTGCGCGGCCAGGTCCCCCGCGGACCTTCACCTGCCAGATGACGTGGCACCCGCG	1060		
Sbjct 1097	CGGCGGTGCGCGGCCAGGGCCCCCGCGCACCTTCACCTGCCAGTGACGTGGCACCCGCG	1156		
Query 1061	ACTCCGTGATGTTCTCGCGACGCAATGCCACCGGGCTGGCCCTGGTGTGCCCGCGCCAA	1120		
Sbjct 1157	ACTCCGTGTGTTCTCTCGGCGCAACGCCAGCGGCACGGCATCGGTGCTGCCCGCGCCAA	1216		
Query 1121	CCATACCATGGAATTTGGGGTCCGGCATGTGGTCTGCACGGCCGGCTGCGTCCCCGAGG	1180		
Sbjct 1217	CCATTACCATGGAGTTTACGGGCGACCATGCGGTCTGCACGGCCGGCTGTGTGCCCGAGG	1276		
Query 1181	GCGTGACGTTT~CCTGGTTCCCTGGGGGACGACCCCTCACCGGCGGCTAAGTCGGCCGTTA	1240		
Sbjct 1277	GGGTGACGTTTGCTGCTGCTTCTGGGGGACGACTCCTCGCCGGCGGAGAGGTGGCCGTCG	1336		
Query 1241	CGGCCCAGGAGTCGTGCGACACCCCGGGCTGGCTACGGTCCGGTCCACCTGCCCATTT	1300		
Sbjct 1337	CGTCCCAGACATCGTGCGGGCGCCCCGGCACCGCCACGATCCGCTCCACCTGCCGCTCT	1396		
Query 1301	CGTACGACTACAGCGAGTACATCTGTCGGTTGACCGGATATCCGGCCGGGATTCCCGTTC	1360		
Sbjct 1397	CGTACGAGCAGACCGAGTACATCTGCCGGCTGGCGGGATACCCGGACGGAATTCGGTCC	1456		
Query 1361	TAGAGCACCACGGCAGTCACGACGCCCCACCCAGGGACCCACCGAGCTGCAGGTGATCG	1420		
Sbjct 1457	TAGAGCACCACGGCAGCCACGACGCCCCCGCGGGGACCCACCGAGCGGCAGGTGATCC	1516		
Query 1421	AGGCGATCGATTGGGTGGGGATTGGA 1446			
Sbjct 1517	GGGCGGTGGAGGGGGCGGGGATCGGA 1542			

Detection of Amino acid sequences of Glycoprotein C and F

The amino acid sequences of both Glycoprotein C and F were detected by Protein data bank PDB as indicated below

Herpes simplex virus type 1 glycoprotein C

translation="MAPGRVGLAVVLSLLWLGAGVSGGSETASTGPT
ITAGAVTNASEAPTSGSPGSAASPEVTPSTPNPNNVTQNKTP
TEPASPTTPKPTSTPKSPPTSTPDPKPKNNTPAKSGRPTKPP
GPVWCDRRDPLARYGSRVQIRCFRNSTRMEFRLQIWRYSMGP
SPPIAPAPDLEEVLNITAPPGGLLVYDSAPNLTDPHVLWAEGAG
PGADPLYSVTGPLPTQRLIIGEVTPATQGMYYLAWGRMDSPHE
YGTWVRVRFMRPPSLTLQPHAVMEGQPFKATCTADAYYPRNPV
ELVWFEDDRQVFNPQGIDTQTHEHPDGFTTVSTVTSEAVGGQVP
PRTFTCQMTWHRDSVTFSSRNATGLALVLRPTITMEFGVRHV
CTAGCVPEGVTFWFLGDDPSAASKAVTAQESCDHPGLATVRST
LPISYDYSEYICRLTGYPAGIPVLEHHGSHQPPRPDTERQVIEAIE
WVGIGIGVLAAGVLVVTIAIVYVVRTSQSRQRHRR"

Herpes simplex virus type 2 glycoprotein C

/translation="MAFRASGPAYQPLAPRPPPARARVPAVAWIGVGAI

VGAFALVAALVLVPPRSSWGLCPDSGWQEFNAGCVAWDPTPVE
HEQAVGGCSAPATLIPRAAAKHLAALTRVQAERSSGYWWVNGDGI
RTCLRLVDSVSGIDEFCEELMALGRVGLTVGLWGLLVGVVVVLA
NASPG RTITVGPGRNASNAAPSVPRNRSAPRTTPTPPQPRKATK
SKASTAKPAPPPKTGPPKTSSEPVRCNRHDPLARYGS
RVQIRCFPNSTRTESRLQIWRYATATDAEIGTAPSLEEVMNVSA
PPGGQLVYDSAPNRTDPHVIWAEGAGPGASPRLYSVVGLGRQRL
HIEELTLETQGMYYVWVWGRTRPSAYGTWVRVRFPRPSLTIHPH
AVLEGQPFKATCTAATYYPGNRAEFVWFEDGRRVFDPAQIHTQT
QENPDGFSTVSTVTSAAVGGQGPRTFTCQLTWHRDSVFSRRN
ASGTASVLRPTITMEFTGDHAVCTAGCVPEGVTFWFLGDDSS
PAEKVAVASQTSCGRPGTATIRSTLPVSYEQTEYICRLAGYPDGIPV
LEHHGSHQPPRPDTERQVIRAVEGAGIGVAVLVAVVLAGTAVVYL
THASSVRYRRLR

Alignment between glycoprotein C and other strains and isolates

The alignment between glycoprotein C and F of Herpes simplex virus type 1 and 2 respectively and some other strains and isolates indicates around (99-100%) homology.

Prediction of the structure of Glycoprotein C and F (Table 3 & 4)

Table 3: Alignment between glycoprotein C and other strains and isolates.

1.Human herpesvirus 1 isolate E5 glycoprotein C (UL44) gene, complete cds (100%)				
Score	Expect	Identities	Gaps	Strand
2837 bits (1536)	0	1536/1536(100%)	0/1536(0%)	Plus/Plus
2. Human herpesvirus 1 strain KOS, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
2793 bits (1512)	0	1528/1536(99%)	0/1536(0%)	Plus/Plus
3. Human herpesvirus 1 strain KOS63, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
2793 bits (1512)	0	1528/1536(99%)	0/1536(0%)	Plus/Plus
4. Human alpha herpesvirus 1 strain 2015-30637, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
2787 bits (1509)	0	1527/1536(99%)	0/1536(0%)	Plus/Plus

Table 4: Alignment between Glycoprotein F and other strains and isolates.

1. Human alpha herpesvirus 2 strain 2008_83, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
4170 bits (2258)	0	2285/2297(99%)	5/2297(0%)	Plus/Plus
2. Human alpha herpesvirus 2 strain 2012_3154, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
4170 bits (2258)	0	2285/2297(99%)	5/2297(0%)	Plus/Plus
3. Human alpha herpesvirus 2 strain 2001_6349, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
4170 bits (2258)	0	2285/2297(99%)	5/2297(0%)	Plus/Plus

4. Human alpha herpesvirus 2 strain 2015_14086, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
4165 bits (2255)	0	2284/2297(99%)	5/2297(0%)	Plus/Plus
5. Human alpha herpesvirus 2 strain 2014_23324, partial genome (99%)				
Score	Expect	Identities	Gaps	Strand
4165 bits (2255)	0	2284/2297(99%)	5/2297(0%)	Plus/Plus

Structure of glycoprotein C: The below structure of glycoprotein C presented by PDB indicates tertiary (three dimensional) structure that harbors both α helices and β pleated sheets (Figure 1)

Structure of glycoprotein F: The above structure of glycoprotein F presented by PDB indicates tertiary (three dimensional) structure that harbors both α helices and β pleated sheets (Figure 2).

Phyre2 to analyze protein structure

Analysis of the structure of glycoprotein C and its functions

(Figure 3): Phyre2 is a suite of tools available on the web to predict and analyze protein structure, function and mutations. It showed the followings:

Figure 3: Analysis of the structure of glycoprotein C and its functions.

- a. The complex crystal structure of glycoprotein C

b. The antigenic determinant site that binds with the fab region of specific immunoglobulin

c. The cytokine receptor site

d. The structural basis of interferon

e. The cell adhesion site
- Phyre2 showed the followings (Figure 4):

a) The heterodimer structure of glycoprotein F

b) The antigenic determinant site that binds with the fc fragment region of specific immunoglobulin

c) Solution structure of human secretory IgA

d) Enzyme deglycosylase human IgG Fc fragment

e) The complex crystal structure of glycoprotein F

Figure 4: Analysis of the structure of glycoprotein F and its functions.

Discussion

The recent flood of data from genome sequences and functional genomics has given rise to new field, bioinformatics, which combines elements of biology and computer science.

Bioinformatics is conceptualizing biology in terms of macromolecules (in the sense of physical-chemistry) and then applying "informatics" techniques (derived from disciplines such as applied math's, computer science, and statistics) to understand and organize the information associated with these molecules, on a large-scale.

We are using Bioinformatics to study homology between the Herpes Simplex virus type 1 glycoprotein C gene and Herpes Simplex virus type 2 glycoprotein F and We found 81% homology, while [1] found the overall sequence homology between these two fragments was 68%. However, certain regions of the sequence showed either a much higher or lower degree of sequence homology

than others. For example, the sequences between positions 0 and 570 of the HSV-1 and HSV-2 sequences showed only 51% homology, whereas the region between positions 570 and 1740 showed a much higher degree of sequence homology (80%). An additional highly homologous region (70%) was also found at the end of the two sequences from position 1975 to position 2419. In addition to the nucleotide sequence changes, the two genomes showed various deletions or insertions when compared with each other. The most notable was an 81-base-pair region found at positions 346 to 426 of the HSV-1 gC sequence which is missing from the HSV-2 genome. From this overall sequence comparison, it appeared that there was a high degree of sequence homology between the HSV-1 region and the HSV-2 gF region sequenced here.

Our results help explain previous results which demonstrated that the HSV-2 gF and HSV-1 gC proteins were mainly type specific but did have type-common determinants [36,38,40,42]. Since several previous studies [32,36] demonstrated that these proteins

induced predominantly type-specific antibodies, it is reasonable that the most antigenic regions of the proteins are found within the more divergent N-terminal sequences which follow the putative hydrophobic signal sequences. The hydrophilic nature of the divergent regions, along with their high content of potential-linked glycosylation sites [45], suggests that these regions would be located on the surface of the protein. Exposure of these divergent sequences to the outside of the proteins maybe responsible for the generation of type-specific antibodies directed against these divergent epitopes. However, type common antibodies could likely also be generated by the more highly conserved carboxy-terminal three-fourths of the proteins, since hydrophilic regions conserved between gC and gF could be exposed to the outside of the proteins and may be, in one case, glycosylated (residues 363 to 366 of HSV-1 gC and 330 to 332 of HSV-2 gF). Thus, HSV-1 gC and HSV-2 gF share both type-specific and type-common determinants, but it appears that the type-specific determinants are more antigenic. Although an explanation of the type-specific and type common determinants of gC and gF must at this point be speculative, it is possible that the proteins have at least two functions, one of which is important for the viability of both viruses, the type-common domain, and one of which is specific for each virus type, the type-specific domain. Although the function(s) of gC and gF is at present unknown and viable gC minus mutants of HSV-1 have been isolated invitro [44], it is not clear that either gC or gF is indispensable to the viruses during in vivo infection of the human host and the establishment of latency. It is possible that at least some of the biological differences between HSV-1 and HSV-2, including prediction for site of infection and virulence, maybe due to the marked structural differences between the amino-terminal regions of gC and gF.

Conclusion

The results reported in this paper demonstrate that the HSV-1 gC and HSV-2 gF glycoproteins are highly homologous and that they encode type-common and type-specific domains.

References

- Donald J. Dowbenko, Laurence A Lasky (1984) Extensive Homology Between the Herpes Simplex Virus Type 2 Glycoprotein F Gene and the Herpes Simplex Virus Type 1 Glycoprotein C Gene. *J Virol* 52(1): 154-163.
- Shan Ling Hung, Sudha Srinivasan, et al. (1992) Structural Basis of C3b Binding by Glycoprotein C of Herpes Simplex Virus. *J Virol* 66(7): 4013-4027.
- Baucke R, P Spear (1979) Membrane proteins specified by herpes simplex viruses. V. Identification of an Fc-binding glycoprotein. *J Virol* 32(3): 779-789.
- Bell S, M Cranage, L Borysiewicz, T Minson (1990) Induction of immunoglobulin G Fc receptors by recombinant vaccinia viruses expressing glycoproteins E and I of herpes simplex virus type 1. *J Virol* 64(5): 2181-2186.
- Draper KG, RH Costa, GTY Lee, P G Spear, E K Wagner (1984) Molecular basis of the glycoprotein-C-negative phenotype of herpes simplex virus type 1 macroplaque strain. *J Virol* 51(3): 578-585.
- Dubin G, I Frank, HM Friedman (1990) Herpes simplex virus type 1 encodes two Fc receptors which have different binding characteristics for monomeric immunoglobulin G (IgG) and IgG complexes. *J Virol* 64(6): 2725-2731.
- Eisenberg RJ, M Ponce de Leon, HM Friedman, LF Fries, MM Frank, et al. (1987) Complement component C3b binds directly to purified glycoprotein C of herpes simplex virus types 1 and 2. *Microb Pathog* 3(6): 423-435.
- Friedman HM, GH Cohen, RJ Eisenberg, CA Seidel, DB Cines (1984) Glycoprotein C of herpes simplex virus 1 acts as a receptor for the C3b complement component on infected cells. *Nature (London)* 309(5969): 633-635.
- Ghosh Choudhury, NM Butcher, HP Ghosh (1990) Expression from cloned DNA of biologically active glycoprotein C of herpes simplex virus type 1 in mammalian cells. *J Gen Virol* 71(Pt 3): 689-699.
- Hanke T, FL Graham, V Lulitanand, DC Johnson (1990) Herpes simplex virus IgG Fc receptors induced using recombinant adenovirus vectors expressing glycoproteins E and I. *Virology* 177: 437-444.
- Herold BC, D WuDunn, N Soltys, PG Spear (1991) Glycoprotein C of herpes simplex virus type 1 plays a principal role in the adsorption of virus to cells and in infectivity. *J Virol* 65(3): 1090-1098.
- Holland TC, FL Homa, SD Marlin, M Levine, J Glorioso (1984) Herpes simplex virus type 1 glycoprotein C-negative mutants exhibit multiple phenotypes, including secretion of truncated glycoproteins. *J Virol* 52(2): 566-574.
- Hourcade D, VM Holers, JP Atkinson (1989) The regulators of complement activation (RCA) gene cluster. *Adv Immunol* 45:381-416.
- Huemer HP, M Broker, C Larcher, JD Lambris, MP Dierich (1989) The central segment of herpes simplex virus type 1 glycoprotein C (gC) is not involved in C3b binding: demonstration by using monoclonal antibodies and recombinant gC expressed in *Escherichia coli*. *J Gen Virol* 70: 1571-1578.
- Johnson DC, V Feenstra (1987) Identification of a novel herpes simplex virus type 1-induced glycoprotein which complexes with gE and binds immunoglobulin. *J Virol* 61(7): 2208-2216.
- Johnson DC, MC Frame, MW Ligas, AM Cross, ND Stow (1988) Herpes simplex virus immunoglobulin G Fc receptor activity depends on a complex of two viral glycoproteins, gE and gI. *J Virol* 62(4): 1347-1354.
- Johnson DC, MR McDermott, C Chrisp, JC Glorioso (1986) Pathogenicity in mice of herpes simplex virus type 2 mutants unable to express glycoprotein C. *J Virol* 58: 36-42.
- Klickstein, LB, WW Wong, JA Smith, JH Weis, JG Wilson et al. (1987) Demonstration of long homologous repeating domains that are composed of the short consensus repeats characteristic of C3/C4 binding proteins. *J Exp Med* 165(4): 1095-1112.
- Krych M, D Hourcade, JP Atkinson (1991) Sites within the complement C3b/C4b receptor important for the specificity of ligand binding. *Proc Natl Acad Sci USA* 88(10): 4353-4357.
- Kubota Y, TA Gaither, J Cason, JJ O'Shea, TJ Lawley (1987) Characterization of the C3 receptor induced by herpes simplex virus type 1 infection of human epidermal, endothelial, and A431 cells. *J Immunol* 138(4): 1137-1142.
- McNearney TA, C Odell, VM Holers, PG Spear, JP Atkinson (1987) Herpes simplex virus glycoproteins gC-1 and gC-2 bind to the third component of complement and provide protection against complement-mediated neutralization of viral infectivity. *J Exp Med* 166(5): 1525-1535.
- Para MF, RB Baucke, PG Spear (1980) Immunoglobulin G (Fc)-binding receptors on virions of herpes simplex virus type 1 and transfer of these receptors to the cell surface by infection. *J Virol* 34(2): 512-520.
- Perkins SJ, PI Haris, RB Sim, D Chapman (1988) A study of the structure of human complement component factor H by Fourier transform infrared spectroscopy and secondary structure averaging methods. *Biochemistry* 27(11): 4004-4012.
- Reid KB, DR Bentley, RD Campbell, LP Chyng, RB Sim, et al. (1986) Complement system proteins which interact with C3b or C4b. *Immunol Today* 7(7-8): 230-234.

25. Sears AE, BS McGwire, B Roizman (1991) Infection of polarized MDCK cells with herpes simplex virus 1: two asymmetrically distributed cell receptors interact with different viral proteins. *Proc Natl Acad Sci USA* 88(12): 5087-5091.
26. Seidel-Dugan C, M Ponce de Leon, HM Friedman, RJ Eisenberg, GH Cohen (1990) Identification of C3b-binding regions on herpes simplex virus type 2 glycoprotein C. *J Virol* 64(5): 1897-1906.
27. Seidel-Dugan C, M Ponce de Leon, HM Friedman, LF Fries, MM Frank et al. (1988) C3b receptor activity on transfected cells expressing glycoprotein C of herpes simplex virus types 1 and 2. *J Virol* 62: 4027-4036.
28. Tal-Singer R, C Seidel-Dugan, L Fries, HP Huemer, RJ Eisenberg et al. (1991) Herpes simplex virus glycoprotein C is a receptor for complement component iC3b. *J Infect Dis* 164(4): 750-753.
29. Zezulak KM, PG Spear (1984) Mapping of the structural gene for the herpes simplex virus type 2 counterpart of herpes simplex virus type 1 glycoprotein C and identification of a type 2 mutant which does not express this glycoprotein. *J Virol* 49(30029):741-747.
30. Balachandran N, D Harnish, RA Killington, S Bacchetti, WE Rawls (1981) Monoclonal antibodies to two glycoproteins of herpes simplex virus type 2. *J Virol* 39(2): 438-446.
31. Balachandran N, D Harnish, WE Rawls, S Bacchetti (1982) Glycoproteins of herpes simplex virus type 2 as defined by monoclonal antibodies. *J Virol* 44: 344-355.
32. Eberle R, RJ Courtney (1981) Assay for type-specific and type-common antibodies to herpes simplex virus types 1 and 2 in human sera. *Infect Immun* 31(3): 1062-1070.
33. Eisenberg RJ, M Ponce de Leon, L Pereira, D Long, GH Cohen (1982) Purification of glycoprotein gD of herpes simplex virus types 1 and 2 by use of monoclonal antibody. *J Virol* 41(3): 1099-1104.
34. Norrild B, H Ludwig, R Rott (1978) Identification of a common antigen of herpes simplex virus, bovine herpes mammillitis virus, and B virus. *J Virol* 26(3): 712-717.
35. Para MF, L Goldstein, PG Spear (1982) Similarities and differences in the Fc-binding glycoprotein (gE) of herpes simplex virus types 1 and 2 and tentative mapping of the viral gene for this glycoprotein. *J Virol* 41(1): 137-144.
36. Pereira L, D Dondero, B Norrild, B Roizman (1981) Differential immunologic reactivity and processing of glycoproteins A and gB of herpes simplex virus types 1 and 2 made in Vero and HEP-2 cells. *Proc Natl Acad Sci USA* 78(8): 5202-5206.
37. Pereira LT, Klassen T, JR Baringer (1980) Type-common and type-specific monoclonal antibody to herpes simplex virus type 1. *Infect Immune* 29(2): 724-732.
38. Powell K, A Buchan, C Sim, D Watson (1974) Type specific protein in herpes simplex virus envelope reacts with neutralizing antibody. *Nature (London)* 249(455): 360-361.
39. Spear P (1980) Herpes viruses. Cell membranes and viral envelopes. HA Blough, JM Tiffaney (ed) Academic Press Inc London 2: 709-750.
40. Zezulak KM, PG Spear (1983) Characterization of a herpes simplex virus type 2 75,000-molecular-weight glycoprotein antigenically related to herpes simplex virus type 1 glycoprotein C. *J Virol* 47(3): 553-562.
41. Zezulak KM, PG Spear (1984) Mapping of the structural gene for the herpes simplex virus type 2 counterpart of herpes simplex virus type 1 glycoprotein C and identification of a type 2 mutant which does not express this glycoprotein. *J Virol* 49(3): 741-747.
42. Zweig M, SD Showalter, SV Bladen, CJ Heilman Jr, B Hampar (1983) Herpes simplex virus type 2 glycoprotein F and type 1 glycoprotein gC have related antigenic determinants. *J Virol* 47(1): 185-192.
43. Homa FL, DJ Purifoy, JC Glorioso, M Levine (1986) Molecular basis of the glycoprotein C-negative phenotypes of herpes simplex virus type 1 mutants selected with a virus-neutralizing monoclonal antibody. *J Virol* 58(2): 281-289.
44. Cassai E, R Manservigi, A Corallini, M Terni (1975) Plaque dissociation of herpes simplex viruses: biochemical and biological characters of viral variants. *Intervirology* 6(4-5): 212-223.
45. Hubbard S, R Ivatt (1981) Synthesis and processing of asparagine-linked oligosaccharides. *Annu Rev Biochem* 50: 555-583.