



Case Report

Copy Right@ Stefan Bittmann

# Successful Treatment of a Critical ill Newborn with COVID-19 Pneumonia after Cesarean Section

Fuyong Jiao<sup>1</sup>, Chongchen Zhou<sup>3</sup>, Yibo Cheng<sup>3</sup>, Shuying Luo<sup>3</sup>, Lei Ma<sup>1</sup>, Xinggang Zhen<sup>3</sup>, Yingjun Feng<sup>3</sup>, Min Li<sup>3</sup>, Stefan Bittmann<sup>2\*</sup>

<sup>1</sup>Children's Hospital, Xiatong University, China

<sup>2</sup>Department of Pediatrics, Ped Mind Institute, Germany

<sup>3</sup>Henan Provincial Children's Hospital, Children's Hospital affiliated to Zhengzhou University, China

\*Corresponding author: Stefan Bittmann, Head of the Department of Pediatrics and Ped Mind Institute D-48599 Gronau, Germany.

To Cite This Article: Stefan Bittmann. Successful Treatment of a Critical ill Newborn with COVID-19 Pneumonia after Cesarean Section. 2020 - 9(3). AJBSR.MS.ID.001383. DOI: 10.34297/AJBSR.2020.09.001383.

Received: 📅 June 08, 2020; Published: 📅 June 25, 2020

## Introduction

We report about a 5 day old newborn with critically pneumonia SARS-CoV-2 after cesarean section infected by a family cluster infection. A male newborn was admitted 22 days after delivery due to "intermittent fever for 17 days". That means that on the fifth day after delivery, due to Cesarean section, the newborn began to fever. In short, history of pregnancy and delivery modus was G1P1, 22 days ago (31 Jan 2020). The newborn was delivered in 38 + 5 weeks gestation by c-section. Amniotic fluid, umbilical cord and placenta was normal, APGAR score at 1, 5 and 10 minutes normal. 18 days ago (04 Feb 2020) the child developed fever with a peak of 38.0 °C. The child showed choking cough, had no running nose, no vomiting, no diarrhoea, any convulsions or skin rashes.

17 days ago, on 02 of February 2020, the fever in outpatient clinic in Xinyang Center Hospital processed. A swab test for SARS-CoV-2 virus nucleic acid testing was positive. 15 days ago, on 6th of February 2020, fever was present once up to 37.7 °C. 17 days before the child see a doctor in the center of the Xinyang Hospital. The medical treatment was based on ribavirin, cephalosporin, immunoglobulin (2.5 g, application 5 days), albumin (3 g, application of 3 days), atomization interferon, Lopinavir/Ritonavir (0.5 ml /per time, all 12h ). Under these medications the child had temperature fluctuations with highest temperature of 37.5 °C, discontinuous, when administering oxygen inhalation through nasal catheter oxygen intolerance. The child had no failure to thrive, but had choking cough. For further treatment, with a diagnosis of "COVID-19" the newborn was admitted to our hospital from emergency ambulance. In this moment of admittance, the child's mental reaction was poor, but sleep was normal, the newborn showed normal stool and normal urine.

The parents had been living in Wuhan City, before the baby was born. On January 22, the parents of the baby returned to Xinyang City from Wuhan. In the afternoon of that day, the mother felt cold and drowsy. The father and grandfather of the child were diagnosed with COVID-19 on April 4th. The baby's grandmother was diagnosed on February 25. Physical examination showed full month appearance, weight of 3600g, body temperature was 36.5°C, pulse rate 172 times/min, respiration rate of 40 times/min. The blood pressure was initially measured by 74/36mm Hg, height was 51cm. There was no malformation in the skull, the anterior fontanelle was 2.5cm × 2.5cm, the posterior fontanelle was closed, flat and soft, and the tension was normal. Bilateral pupil showed isocorie, the reflex to light existed. The breath was smooth under oxygen inhalation by nasal catheter was performed. No expiratory dyspnoea was found. The three concave signs were negative.

The breath sounds in both lungs were rough. Heart rhythm was uniform, heart sounds were strong, no murmur was heard, abdomen was soft, umbilical cord did not fall off, liver reached 1.5cm below right costal, soft, spleen was not reached, bowel sounds were normal. The extremities were warm at the end, with free movement, normal muscle tone, and primitive reflexes. Laboratory blood routine examination showed leukocyte 12.95×10<sup>9</sup>/L, erythrocyte 3.07×10<sup>12</sup>/L, hemoglobin 101g/L, platelet 356×10<sup>9</sup>/L, lymphocyte percentage 49.5 %, c-reactive protein quantitative determination <0.5mg/L. Liver and kidney function, myocardial enzymes, electrolytes: total bilirubin 13.7 umol/L, conjugated bilirubin 2.7 umol/L, unconjugated bilirubin 11.0 umol/L, alanine transaminase 46.7 U/L, glutamate transaminase 37.8 U/L, creatine kinase isoenzyme 48.3 U/L, potassium 6.75 mmol/L (Figure 1).



