



Opinion

Copy Right@ Juan Pedro Lapuente

CROSSTALK to Fight against Inflammation and Fibrosis: New Biological Medical Drugs Design for a New Age

Juan Pedro Lapuente*

R4T Molecular and Cell Biology research Laboratories, Fuenlabrada University Hospital, Spain.

***Corresponding author:** Juan Pedro Lapuente, Molecular and cell Biology SCO of R4T Molecular and Cell Biology research Laboratories, Fuenlabrada University Hospital, Madrid 28942, Spain.

To Cite This Article: Juan Pedro Lapuente. *CROSSTALK to Fight against Inflammation and Fibrosis: New Biological Medical Drugs Design for a New Age.* *Am J Biomed Sci & Res.* 2022 - 16(1). *AJBSR.MS.ID.002192.* DOI: [10.34297/AJBSR.2022.16.002192](https://doi.org/10.34297/AJBSR.2022.16.002192)

Received: 📅 April 08, 2022; **Published:** 📅 April 14, 2022

Opinion

Inflammation is the response of an organism's immune system to damage caused to its cells and vascularized tissues by any type of aggression, which may be of a biological, chemical, or physical nature. This inflammatory response must be self-limiting in time and intensity, since, if this is not the case and there is not perfect coordination between the innate and adaptive immune systems, at the very least a fibrotic process will be generated that will lead to a percentage loss of function in the affected tissue or tissues, and, in severe cases, could lead to a severe systemic inflammatory syndrome with positive feed-back systems, leading to a cytokine storm, which in turn could lead to multi-organ failure [1,2]. In the establishment, maintenance and termination of such a cytokine storm at the molecular level, soluble cytokine-like mediators (especially IL-1, IL-6, IL-18, etc...), growth factors (TNF- α , etc...), chemokines (MCP-1, MIP-1, etc...), molecules related to the cytokines (MCP-1, MIP-1, etc...) and cytokine-related molecules (MCP-1, MIP-1, etc...) are essential for the establishment, maintenance and termination of such a cytokine storm.), molecules related to the purinergic system (ATP, ADP), which will act on receptors such as Toll-like Receptors (TLRs), NOD-like Receptors (NLRs), RIG-type Helicase Receptors (RLRs) and purinergic receptors (P2X7).

Biopharmaceuticals [3], as their name suggests, are manufactured from biological sources, and can be formulated synthetically, semi-synthetically or even completely naturally. The definition of biological medicines encompasses a very

heterogeneous range of medicines, such as whole or partial blood components, cell therapies, tissues, proteins, vaccines, and gene therapies.

Within this group, so-called biologics are becoming increasingly prominent, as they have clear competitive advantages over advanced cell therapy medicines or gene therapy medicines. With respect to advanced cell therapy drugs, biologics are cell-free and have several advantages that allow them to have fewer risks, fewer limitations, and greater efficacy than treatments based on the direct use of cells. For example, biological drugs are not influenced by aspects such as the age or sex of the donor, which is the case with advanced cell therapy drugs [4-8], and biological drugs also carry less risk when injected than drugs that contain cells in their formulation [9]. This, together with aspects related to their manufacture and commercialization, such as wide scalability, storage capacity, fractionation, concentration, and the possibility of combining materials from different donors in their manufacture, make them much more interesting [10,11]. Finally, another aspect to highlight as an advantage of the new biologic drugs over advanced cell therapy is that the manufacturing method of these new biologic drugs can direct the response towards a specific cell type or tissue, achieving a tissue selectivity that is difficult to obtain in cell therapy. We must not forget that stem cells are organized in the organism in specific niches characteristic of each type of tissue and are clearly influenced by the cellular and extracellular microenvironment, so that all the secretome they produce, both



in soluble form (cytokines, etc.) and encapsulated in exosomes (miRNA, etc.), is tissue selective [12,13].

Among biological drugs, the most developed in recent years are monoclonal antibodies, specialized glycoproteins that are part of the immune system, produced by B-lymphocytes, with the ability to recognize specific molecules (antigens). Monoclonal antibodies have become essential clinical and biotechnological tools and have proven useful in the diagnosis and treatment of infectious, immunological and neoplastic diseases, but although these recombinant monoclonal antibodies are currently manufactured using phage library technology with genes encoding variable regions of immunoglobulins, These antibodies are not free from problems related mainly to immunogenicity reactions, which ultimately lead to variability in their efficacy, rendering them ineffective in a significant percentage of the patients to whom they are applied [14,15]. However, these losses in efficacy are not only due to the existence of immunogenicity reactions, including the generation of anti-drug antibodies, but also to the special mode of action that derives from their name, since monoclonal antibodies act against a single, specific target.

It is very difficult to explain the existence of a cytokine storm by the activation of a single receptor. If this were the case, treatment of the cytokine storm by a single monoclonal antibody, e.g., a monoclonal antibody against IL-1 β or IL-6, would always be effective, and we know that this is almost never the case, as the SARS-COV-2 pandemic has recently demonstrated. Moreover, even if only one of the receptors is activated in the first instance, the simple start-up of its metabolic cascade will provoke the appearance of DAMPS (Self Damaging Molecular Patterns) that will stimulate other receptors. If we add to this the fact that in most cytokine storms, we do not see a single causative agent, but rather a group of them, we can understand that there is almost always a joint activation of several of these receptors, producing phenomena of agonism, synergy and antagonism between them [16]. Finally, if all this inflammatory tsunami is not controlled in time, the dreaded fibrosis with the consequent loss of function of the affected tissue, and even, in severe cases, autoimmune reactions, will appear as a minimum.

But in the cytokine storm we must also consider the involvement of the purinergic system [17-19]. Extracellular ATP or its enzymatic degradation products, such as ADP, AMP, and adenosine, can stimulate several membrane receptors [20]. Stimulation of the P2X7 receptor by ATP leads to activation of the NLRP3 inflammasome, and consequently of caspase 1, stimulating exaggerated secretion of IL-1 β and IL-18 [21].

Based on the above, it is logical to think that to produce a global anti-inflammatory immunomodulatory effect that acts synergistically at all levels, avoiding feedback pathways that prolong inflammation over time with all its negative consequences, complex biological drugs are needed that act at multiple levels, in a synergistic and coordinated manner, avoiding the negative consequences of blocking, or stimulating a single pathway. For all these reasons, our research group at the R4T molecular and cellular biology research laboratory of the Hospital Universitario de Fuenlabrada, unlike the treatments tested to date to control cytokine storms based on the use of monoclonal antibodies used alone or in combination, proposes a biological therapy based on the use of allogeneic conditioned medium derived from M2-type macrophages and enriched with MSCs as a treatment.

Mesenchymal stem cells, placed in coculture with macrophages, not only respond to macrophages and adjust their secretome accordingly, but also macrophages respond to encounters with these cells, creating a feedback loop that contributes to immune regulation [22]. Our technological platform for the manufacture of complex biological drugs based on the establishment of an indirect cellular crosstalk allows us to generate various conditioned media, where all the growth factors, cytokines, chemokines, exosomes, miRNA, etc., that are naturally produced by these cells are present, respecting their composition and natural pleiotropic relationships, with an immunomodulatory cytokine profile that confers a powerful anti-inflammatory and antifibrotic action, in a totally tissue-selective manner.

For all these reasons, we are firmly convinced of the advantages of using complete conditioned media, manufactured through technological platforms that allow intercellular crosstalk between two or more specific cell types, as opposed to one of their purified components, given that their efficacy lies in the synergistic mechanism between their different components [23], the result of subjecting cell populations to a culture that, *in vitro*, attempts to emulate the anti-inflammatory, anti-fibrotic and regenerative immunomodulatory microenvironment that should occur under ideal *in vivo* conditions in diseased tissue.

References

1. Zindel J, Kubes P (2020) DAMPs, PAMPs, and LAMPs in Immunity and Sterile Inflammation. *Annu Rev Pathol* 15: 493-518.
2. Medzhitov R (2008) Origin and physiological roles of inflammation. *Nature* 454(7203): 428-435.
3. Rader RA (2008) (Re)defining biopharmaceutical. *Nat Biotechnol* 26(7): 743-751.
4. Efimenko A, Dzhoyashvili N, Kalinina N, Kochegura T, Akchurin R, et al. (2014) Adipose-derived mesenchymal stromal cells from aged patients

- with coronary artery disease keep mesenchymal stromal cell properties but exhibit characteristics of aging and have impaired angiogenic potential. *Stem Cells Translational Medicine* 3(1): 32-41.
5. Efimenko A, Starostina E, Kalinina N, Stolzing A (2011) Angiogenic properties of aged adipose derived mesenchymal stem cells after hypoxic conditioning. *Journal of Translational Medicine* 9(1): 10.
 6. Mohd Ali N, Boo L, Yeap SK, Ky H, Satharasinghe DA, et al. (2016) Probable impact of age and hypoxia on proliferation and microRNA expression profile of bone marrow derived human mesenchymal stem cells. *Peer J* 14(4): 1536.
 7. Siegel G, Kluba T, Hermanutz Klein U, Bieback K, Northoff H, et al. (2013) Phenotype, donor age and gender affect function of human bone marrow-derived mesenchymal stromal cells. *BMC Medicine* 11(1): 146.
 8. Pasanen I, Pietilä M, Lehtonen S, Lehtilahti E, Hakkarainen T, et al. (2015) Mesenchymal stromal cells from female donors enhance breast cancer cell proliferation in vitro. *Oncology* 88(4): 214-225.
 9. Herberts CA, Kwa MS, Hermsen HP (2011) Risk factors in the development of stem cell therapy. *Journal of Translational Medicine* 9(1): 29.
 10. Nath SC, Harper L, Rancourt DE (2020) Cell-Based Therapy Manufacturing in Stirred Suspension Bioreactor: Thoughts for cGMP Compliance. *Front Bioeng Biotechnol* 8: 599674.
 11. Maitra B, Szekely E, Gjini K, Laughlin MJ, Dennis J, et al. (2004) Human mesenchymal stem cells support unrelated donor hematopoietic stem cells and suppress T-cell activation. *Bone Marrow Transplant* 33(6): 597-604.
 12. Baglio SR, Rooijers K, Koppers Lalic D, Verweij FJ, Pérez Lanzón M, et al. (2015) Human bone marrow- and adipose-mesenchymal stem cells secrete exosomes enriched in distinctive miRNA and tRNA species. *Stem Cell Research & Therapy* 6(1): 127.
 13. Strioga M, Viswanathan S, Darinskas A, Slaby O, Michalek J (2012) Same or not the same? Comparison of adipose tissue-derived versus bone marrow-derived mesenchymal stem and stromal cells. *Stem Cells and Development* 21(14): 2724-2752.
 14. López Ibáñez M, Marín Jiménez I (2016) Pharmacy and Anti-Pharmacy Levels in the Clinical Care of the Patient with Intestinal Inflammation [Drugs and anti-drug antibody levels in the management of patients with inflammatory bowel disease]. *Gastroenterol Hepatol* 39(4): 265-272.
 15. Stidham RW, Lee TC, Higgins PD, Deshpande AR, Sussman DA, et al. (2014) Systematic review with network meta-analysis: the efficacy of anti-TNF agents for the treatment of Crohn's disease. *Aliment Pharmacol Ther* 39(12): 1349-1362.
 16. Root Bernstein R (2021) Innate Receptor Activation Patterns Involving TLR and NLR Synergisms in COVID-19, ALI/ARDS and Sepsis Cytokine Storms: A Review and Model Making Novel Predictions and Therapeutic Suggestions. *Int J Mol Sci* 22(4): 2108.
 17. Simões JLB, Bagatini MD (2021) Purinergic Signaling of ATP in COVID-19 Associated Guillain-Barré Syndrome. *J Neuroimmune Pharmacol* 16(1): 48-58.
 18. Franciosi MLM, Lima MDM, Schetinger MRC, Cardoso AM (2021) Possible role of purinergic signaling in COVID-19. *Mol Cell Biochem* 476(8): 2891-2898.
 19. Simões JLB, de Araújo JB, Bagatini MD (2021) Anti-inflammatory Therapy by Cholinergic and Purinergic Modulation in Multiple Sclerosis Associated with SARS-CoV-2 Infection. *Mol Neurobiol* 58(10): 5090-5111.
 20. Abbracchio MP, Burnstock G (1994) Purinoceptors: Are there families of P2X and P2Y purinoceptors? *Pharmacol Ther* 64: 445-475.
 21. Ren W, Rubini P, Tang Y, Engel T, Illes P (2021) Inherent P2X7 Receptors Regulate Macrophage Functions during Inflammatory Diseases. *Int J Mol Sci* 23(1): 232.
 22. Carty F, Mahon BP, English K (2017) The influence of macrophages on mesenchymal stromal cell therapy: passive or aggressive agents? *Clin Exp Immunol* 188(1): 1-11.
 23. Laggner M, Gugereil A, Bachmann C, Hofbauer H, Vorstandlechner V, et al. (2020) Reproducibility of GMP-compliant production of therapeutic stressed peripheral blood mononuclear cell-derived secretome, a novel class of biological medicinal products. *Stem Cell Res Ther* 11(1): 9.