



Research Article

Copy Right@ Saccomanni Bernardino

Lumbar Disc Erniation l4-l5; l5-s1: An Original Study

Saccomanni Bernardino*

Orthopaedic and Trauma Surgery, Asl Bari, Italy

*Corresponding author: Saccomanni Bernardino, Orthopaedic and Trauma Surgery, Asl Bari, Italy.

To Cite This Article: Saccomanni B. Lumbar Disc Erniation l4-l5; l5-s1: An Original Study. Am J Biomed Sci & Res. 2022 17(4) AJBSR.MS.ID.002362, DOI: [10.34297/AJBSR.2022.17.002362](https://doi.org/10.34297/AJBSR.2022.17.002362)

Received: 📅 November 14, 2022; Published: 📅 November 23, 2022

Objectives or Purpose or Study Design

To evaluate the Functional & Neurological outcome of Lumbar Microdiscectomy at L4/5 and L5/S1.

Background: Lumbar disc herniation (LDH) is the most common specific cause for low back pain (LBP). It is a degenerative process causing annular tear with extrusion of the nucleus pulposus through posterior midline or posterolateral aspect of the disc leading to compression of the thecal sac and nerve roots with radicular symptoms. Surgery is reserved for only those who are refractory for a fair trial of nonsurgical management for at least 6 weeks, need of decompression for conus and cauda equina compression, needing immediate attention. There have been various modalities of treatment that have evolved over the years. Micro-lumbar discectomy is one of the latest additions of surgical management modality.

Materials & Method: Retrospectively we reviewed 30 cases from January 2010 to December 2012 in hospital in ALTAMURA ITALY. All the patients had back pain & some neurological deficit with positive MRI findings, underwent lumbar microdiscectomy at L4/5 & L5/S1. Functional improvement was assessed by the ODI Score & categorized into excellent, good, fair & poor, Neurological improvement was assessed by the Nurick Score improvement of pain control by VAS score & overall functional assessment by modified Odom's score.

Results: The total number of patients was 30. Average age was 36 years (range 22-50). The average follow-up period was 9 months. Pre-operative ODI score was 54.6 ± 12.8 & post-operative score was improved to 10.2 ± 3.3 . In our series we have no chair bound or bedridden patients after surgery. The Pre-operative & Postoperative Nurick score was 2.9 & 0.8 respectively. There was no difference in neurological outcomes comparing patients older and younger than 40 years. There was significant improvement in Pre-operative & Post-operative VAS score for leg pain was 6.90 ± 1.9 & 2.0 ± 0.8 respectively & for low back pain was 5.8 ± 1.2 & 2.2 ± 0.8 respectively.

Conclusion: Lumbar Microdiscectomy is a safe & effective method to treat patients with lumbar disc prolapse with shorter hospital stay & better prognosis with minimum & considerable complications.

Keywords: Lumbar Disc; Original Study

Introduction

Backache resulting from Lumbago-sciatica is one of the oldest diseases of history. For decades lumbar disc disease has been a commonly encountered spine problem for spine surgeons. Disc herniation causes radicular pain with or without neuro-deficit and those who do not respond to conservative management for at least 6 weeks, having their activities of daily living affected are managed surgically (Figure 1).

Discectomy for symptomatic lumbar disc herniation is performed commonly performed surgical spinal procedure [1,2]. The first lumbar discectomy was done by a laminectomy and transdermal approach was performed by Miter and Barr in 1934. Semmes first described the hemi-laminectomy approach with retraction of the dura to decompress the neural elements. Discectomy by laminectomy was the common approach for



prolapsed disc for a very long time. However, this involved removal of a large amount of normal bone, ligamentum flava, muscle tissue and sometimes facet joints which resulted in iatrogenic instabilities to the spine and failed back syndromes. Hence, conventional laminectomy and discectomy have been replaced by soft tissue techniques. With the development of better retractor systems, illumination and magnification, discectomies are performed

by a more conservative route of interlaminar approaches. Love described his interlaminar fenestration technique as early as 1939. Surgeons have modified Love's technique to make it minimally invasive [3]. Conventional fenestration technique used paraspinous muscular elevation bilaterally, and larger incisions and retractor systems.




Figure 1: Pathology of Lumbar Disc.

Interlaminar approach was used to enter the epidural space, with aggressive discectomy. Patient selection and/or preoperative evaluation of the patients should be carefully done to increase patient's outcome after the microdiscectomy. In young and athletic patients, we prefer microdiscectomy, because strong paravertebral muscles will share the loading with spinal column. In minimally invasive techniques, paraspinous muscular elevation is done for only 2 to 3 cm using specialised retractor systems. Caspar in 1977 and Williams in 1978 described micro lumbar discectomy technique [4,5]. Due to the postulated advantages of reduced tissue invasiveness, limited blood loss, a shorter duration of surgery, and a faster postoperative recovery, minimally invasive microdiscectomy is very popular now a days. Adequate illumination and magnification of the operative field are achieved by the use

of microscopes, operating loupes and head lamps or endoscopes. Minimally invasive techniques have the overall advantage of less tissue scarring and better visualization of the dura, roots and disc spaces and hence are expected to have better postoperative outcomes [6]. The added advantage of this procedure is minimum tissue damage & less chance of fibrosis. Perineural and epidural fibrosis were reported in the subsequent study. Epidural fibrosis/perineural fibrosis is a nightmare to any managing physician and a major bane for the affected patient. There have been various modalities of treatment for this disease condition that have evolved over the years. The objective of our study is to analyze safety, efficacy, and complications following Micro-lumbar discectomy (Figure 2 & 3).




Figure 2: Sagittal MRI of L/S Spine.




Figure 3: Axial MRI of L/S Spine with pilot film.

Materials & Method

Retrospectively we reviewed 30 cases from 2010 to 2012 of Lumbar disc herniation. All patients were presented with the various symptoms and signs with the positive magnetic resonance imaging (MRI) and underwent decompression by micro lumbar discectomy. Functional improvement was assessed by the ODI Score, Neurological improvement was assessed by the Nurick Score, improvement of pain control by VAS score & overall functional assessment done by modified Odom's score. IBMSPSS V26 software was used for statistical analysis.

Selection of the patients for surgery

Inclusion Criteria

Patients with low back pain who are diagnosed as PLID at L4/5 &/or L5/S1 & not responding to conservative treatment or progressive neurological deficit.

Exclusion Criteria

Patients with lumbar disc prolapse other than L4/5 &/or L5/S1, malignancy, tuberculosis & patients with severe co-morbidity for which he/she is unfit for surgery.

Operative Procedure

The patient should have prone position. Careful inspection should Functional & Neurological outcome of Lumbar Microdiscectomy at L4/5 and L5/S1 be done to the eyes, ulnar nerves, and genitalia for the males, and breast for the females to

ensure that excessive pressure does not exist. Abdominal viscera and vessels should be checked to hang free. The surgical area is propped with an antiseptic solution and covered with sterile clothes. The discectomy level should be identified with C-arm before the operation and incision line is marked.

The length of the incision line changes between 1.5 -2 cm, the lower point of incision should be the upper point of lower spinal process. While performing incision, subcutaneous tissue should not be destroyed to avoid fat tissue necroses. Fascia should be opened just lateral border of spinous process to keep supraspinous ligament intact.

After dissection of the muscle tissue on the spinous process and lamina, it should be cared to the two important points. First of them is to save the capsular ligament and second is to leave intact

the interspinous ligament: therefore, retractor should not be forced against interspinous ligament. The next step is to save ligamentum flavum. In L5-S1 level, the ligamentum flavum is opened from the medial side to lateral as a flap like C and fixed with a spinal needle to the lateral wall. Under the microscope magnification, epidural fat tissue retracted medially with nerve root. This procedure should be performed gently, because there is a very thin layer to keep the fat tissue. If this layer is opened, the dispersed fat tissue can prevent seeing the nerve root. The thin layer should be opened just under the nerve root. Some epidural veins can be seen and coagulated with bipolar. After these procedures, disc annulus can easily be found under the nerve root and discectomy is performed. After the discectomy, ligamentum flavum is placed on the epidural fat tissue. The fascia is sutured, and skin is closed with subcutaneous intradermal sutures (Figure 4 & 5).




Figure 4: Percutaneous identification of L5/S1 space by guide pin with C-Arm image. Brain




Figure 5: Per-operative picture.

Results

The retrospective analytical study includes 30 patients who fulfilled the inclusion criteria, were operated by micro lumbar

discectomy for PLID at L4/5 &/or L5/S1. All patients were followed up from 6 months to 1 year (average 9 months) post-operatively (Tables 1-6).

Table 1: Demographic Distribution of Patients (n=30).

Gender (Male/Female)	Age (years)	Hospital stay (days)	Surgery time (min)	Blood loss (ml)	Follow-up (months)
17/13	40.3±10.1	3.60±1.26	32.80±10.47	28±12.67	15±9; L4/5 17; L5/S1 13

Table 2: Pre-operative and post-operative comparison of pain after 12 months (n=30), According to VAS score.

Score Pre Operative	Final Follow-Up	p value (VAS)	
Back pain	6.90±1.9	(Mean±SD)	1.8±0.8 <0.001
Leg pain	5.8±1.2	(Mean±SD)	2.2±0.8 <0.001

Table 3: Pre-operative and post-operative Nurick Score (n=30).

Operative Nurick
2.9 0.8

Table 4: Pre-operative and post-operative ODI Score.

Score		
	Pre-Operative	Post Operative
ODI	54.6±12.8	10.2±3.3

Table 5: Distribution of patients according to post-operative clinical outcome (n=30).

According to Modified Odom's Criteria (Odom et al, 1958) Comprehensive		
	Frequency (n)	Percentage (%)
Outcome		
Excellent	21	70
Percutaneous identification of L5/S1 space by guide pin		
Good	7	23
with C-Arm image		
Fair	2	7
Poor	0	0

Table 6: Distribution of patients according to post-operative complications.

Complications	No. Of patients
Wound dehiscence	0
Radicular pain	3
Granuloma	0
Wound Infection	1
Seroma	0
Dural leakage	1
Discitis	0

Discussion

Patients treated by micro lumbar discectomy were followed up for the period of 12 months. The overall clinical outcome categorized as excellent, good, fair, and poor according to modified Odom's criteria. For statistical analysis good and excellent were

grouped as satisfactory and fair and poor as unsatisfactory. In this study, age range of the patients were 27-55 years, mean age was 40.3±10.1 years and the male to female ratio was 1.3:1 which are comparable to the study of Fabres A et al. [6]. who found mean age 43.1±13.6 [7].

All 30 (100%) patients had low back pain & radicular pain, motor deficit in 6 (20%) patients, sensory deficit in 5 (16.67%) patients, foot drop in 1 (3%) patient, bowel & bladder involvement in 2 (6.6%). This is also comparable to the study of *Raghu et al.* [7]. All were improved significantly ($P < 0.001$) in postoperative group [8]. Regarding the leg pain control according to VAS was 6.90 ± 1.90 preoperatively and significantly decreased to 1.6 ± 0.8 at the last follow-up ($P < 0.001$). This corresponds to the study of *Jaiswal et al.* [8]. In our series the pre & post-operative Nurick score was 2.9 & 0.8 which was satisfactory for the patients [9].

The mean ODI score was 54.6 ± 12.8 & 10.2 ± 3.3 respectively which is comparable to the study of *Jaiswal et al.* [8]. The differences in the ODI scores were statistically significant between the pre and postoperative follow-ups ($p < 0.001$). ODI scores below 40% were graded as good outcomes (success), whereas higher scores were considered partial or total failures. Post-operative superficial wound infection was in 2 (6.7%) Wound infection was managed conservatively by antibiotics according to culture and sensitivity report, improvement of nutritional status, removal of stitch, regular dressing & secondary wound closure. The average peri-operative blood loss was 28 ± 12.67 ml. Per-operative dural leakage occurred in 1 patient, who was managed by dural repair & placement of a drain tube with early removal of drain. The average post-operative hospital stay was 2-4 days. In our study, 21 (70%) patients got satisfactory results and 7 (23%) patients got good results, in 2 (7%) cases had fair outcome according to the Modified Odom's Criteria. In our series, there was no radiological progression after surgery [10-15].

Conclusion

Lumbar Microdiscectomy is a safe & effective method to treat patients with lumbar disc prolapse with shorter hospital stay & better prognosis with minimum & considerable complications.

Acknowledgment

None.

References

1. Shiju A Majeed, Vikraman CS, Vivek Mathew, Anish TS (2013) Comparison of outcomes between conventional lumbar fenestration discectomy and minimally invasive lumbar discectomy: an observational study with a minimum 2-year follow-up. *J Orthop Surg Res* 8: 1-5.
2. Ozer AF Ali Fahir Ozer MD, Murat Cosar MD PhD.
3. Caspar W (1977) A new surgical procedure for lumbar disc herniation causing less tissue damage through a microsurgical approach. In *Lumbar disc adult hydrocephalus* 74-80.
4. Williams RW (1978) Microlumbar discectomy: a conservative surgical approach to the virgin herniated lumbar disc. *Spine* 3(2): 175-182.
5. Alexander Aichmair, Jerry Y Du, Jennifer Shue, Gisberto Evangelisti, Andrew A Sama (2014) Microdiscectomy for the treatment of lumbar disc herniation: an evaluation of reoperations and long-term outcomes. *Evid Based Spine Care J* (5): 77-86.
6. Fabres A, Escudero N, Schiappacasse R, Cerda J, Salazar C, et al. (2021) Outpatient Lumbar Microdiscectomy: Safe, High Patient Satisfaction and Lower Cost. *Ambulatory Surgery*.
7. Raghu V, Ranade D, Patil A, Gotecha S, Punia P, et al. (2021) A comparative study of conventional microlumbar discectomy and endoscopic lumbar discectomies. *International Surger Journal* 8: 3024-3030.
8. Jaiswal A, Kumar S, Reddy S, Jaiswal P (2019) Feasibility and safety of outpatient lumbar microscopic discectomy in a developing country. *Asian Spine Journal* 3(13): 721.
9. Hoppenfeld S, DeBoer P, Buckley R (2012) *Surgical exposures in orthopaedics: the anatomic approach*. Lippincott Williams & Wilkins.
10. Love JG, Camp JD (1937) Root pain resulting from intraspinal protrusion of intervertebral discs: diagnosis and surgical treatment. *JBJS* 19: 776-804.
11. Magee A, Bhaskar IP, Ilett P, Murphy MA, Wang YY (2016) Clinical and health economic benefits of out-patient lumbar microdiscectomies in Australia. *Australasian Journal of Neuroscience* 26: 15-20.
12. Bhattarai B, Sah SB (2019) A Retrospective Observational Analysis on the Outcome of Microlumbar Discectomy among 70 Consecutive Cases in the Tertiary Care Spine Center in Nepal. *Nepal Journal of Neuroscience* 16: 34-39.
13. Asati S, Jain S, Kundnani VG (2020) Tubular Discectomy Versus Conventional Microdiscectomy for the Treatment of Lumbar Disc Herniation: A Comparative Study. *Journal of Minimally Invasive Spine Surgery and Technique* 5: 51-56.
14. Debono B, Sabatier P, Garnault V, Hamel O, Bousquet P, et al. (2017) Outpatient lumbar microdiscectomy in France: from an economic imperative to a clinical standard-an observational study of 201 cases. *World Neurosurgery* 106: 891-897.
15. Azar FM, Canale ST, Beatty JH (2020) *Campbell's Operative Orthopaedics*, E-Book. Elsevier Health Sciences 23.