



Research Article

Copy Right© : Job Pakisi

Sociodemographic Characteristics and Risk Factors Related to Hypertension Among Individuals from Huambo, Angola

Job Ch Pakisi^{1,2#*}, Cruz S Sebastião^{3,4,5#*}, Jocelyne Vasconcelo³, Leticia Valdeón² and J. Enrique Bayon Darkistade²

¹Faculty of Medicine, José Eduardo dos Santos University, Huambo, Angola

²University of León, León, Spain

³Centro de Investigação em Saúde de Angola (CISA), Caxito, Angola

⁴Instituto Nacional de Investigação em Saúde (INIS), Luanda, Angola

⁵Institute of Health Sciences (ICISA), Agostinho Neto University (UAN), Luanda, Angola

#These authors contributed equally to this work

*Corresponding author: Job Pakisi, Faculty of Medicine, José Eduardo dos Santos University, Huambo, Angola.

Cruz S Sebastião, Centro de Investigação em Saúde de Angola (CISA), Caxito, Angola.

To Cite This Article: Job Pakisi*, Cruz S Sebastião*, Jocelyne Vasconcelo, Leticia Valdeón and Enrique Darkistad, Sociodemographic Characteristics and Risk Factors Related to Hypertension Among Individuals from Huambo, Angola. *Am J Biomed Sci & Res.* 2024 21(3) AJBSR. MS.ID.002849, DOI: [10.34297/AJBSR.2024.21.002849](https://doi.org/10.34297/AJBSR.2024.21.002849)

Received: 📅 January 23, 2024; **Published:** 📅 February 05, 2024

Abstract

WHO estimates that around 600 million people suffer from high blood pressure which causes high costs for health systems and influences the economy of these countries. In this study, we investigate the sociodemographic characteristics and risk factors related to hypertension among individuals from Huambo, Angola. This cross-sectional study was conducted with 158 participants consulted in the Cardiology services at Hospital Geral do Huambo, between January and July 2023. A total of 67.1% of the studied patients were hypertensive. The mean age of the hypertensive (55.8 ± 11.3) patients was higher than non-hypertensive (53.1 ± 13.4) patients, although no significance was observed ($p=0.199$). Patients aged between 30 and 40 ($OR=1.19$, $P=0.493$), male ($OR=1.17$, $P=0.660$), employees ($OR=9.94$, $P=0.935$), patients consuming alcohol ($OR=1.11$, $P=0.864$), and practising physical activity ($OR=1.12$, $P=0.742$) presented a high chance of developing hypertension, while those living in urbanized areas ($OR=0.78$, $P=0.59$) and with a low educational level ($OR=0.43$, $P=0.194$), had a low chance to develop hypertension. There was no statistical significance between demographic or behavioural characteristics with hypertension ($P>0.05$). Regarding clinical characteristics, body mass index was statistically related to hypertension ($p=0.011$). Our findings show that hypertension is a major health problem for the young population of Huambo. There is an urgent need to create strategies to improve the prevention, diagnosis, and treatment of hypertension, not only in Huambo but throughout the country. Further studies to understand determinants related to hypertension should be carried out among the young population in Angola.

Keywords: Hypertension, Risk factors, Huambo, Angola

Introduction

Hypertension is one of the most common chronic diseases, constituting an independent risk factor for many diseases, such as stroke and chronic renal failure [1,2]. The World Health Organization (WHO) estimates that around 600 million people suffer from

high blood pressure, with a global increase of 60% by 2025, in addition to around 7.1 million deaths per year, which causes very high costs for health systems, directly influencing the economy of these countries [3,4]. For example, low- and middle-income countries (LMICs) have 93% of the world's disease burden and contribute to



around 78% of deaths from cardiovascular disease and 50% of the total disease burden [3]. Several authors show that demographic characteristics such as advanced age, alcohol consumption, smoking and a sedentary lifestyle can increase the risk of hypertension [5]. Also, the Body Mass Index (BMI) has been a determining factor in the diagnosis of obesity [5]. Until the 20th century, infectious diseases were the main causes of death in the world population. Today, with the improvement of socio-economic and health conditions, Non-Communicable Diseases (NCDs) are the main cause of death in the world, causing 63% of deaths. In sub-Saharan Africa, the epidemiology of cardiovascular diseases is different compared to other parts of the world, where mortality is declining, and life expectancy is increasing [6].

Studies carried out in Angola on hypertension and associated risk factors, specifically in Bengo, Huambo, Luanda and Lubango, revealed a significant prevalence of hypertension among the young population and a low level of knowledge about this cardiovascular disease [7-10]. Despite that, there are few studies published on the prevalence and risk factors of hypertension in Angola, particularly in Huambo, one of the provinces, located in the centre of the country. Therefore, the present study aimed to investigate the prevalence and factors associated with hypertension in Huambo, in order to contribute to the definition of strategies for controlling cardiovascular diseases.

Material and Methods

Study Design and Setting

This was a cross-sectional study conducted with 158 participants of both genders aged between 30 and 70 years who were consulted at Huambo General Hospital between January and July 2023. Huambo is one of the 18 provinces of Angola, located in the country's central region. According to 2018 population projections, prepared by the National Statistics Institute, it has a population of 2,309,829 inhabitants and a territorial area of 35,771 km², making it the fourth most populous province in Angola and one of the richest in the country. The data were used using the World Health Organization's (WHO) Step Wise questionnaire for surveillance of chronic non-communicable diseases [11]. Sociodemographic (age, gender, occupation, place of residence and educational level), behavioural (alcohol consumption and physical activity) and clinical data (Signs, symptoms and morbidities) were obtained through an interview, after signing the consent form. The research project was first forwarded to the Scientific Council of the Faculty of Medicine of Huambo, having decided positively (Deliberation No. 003/CC/23) and subsequently sent to the Ethics Committee of the Institute of Health Research of Angola, also approved by this Body (031/C.E.M.S./2023). Written informed consent was obtained.

Data and Sample Collection

Blood pressure was measured on the right arm with an OMRON automatic sphygmomanometer with the patient sitting and using the appropriate cuffs according to the individual's upper arm circumference [12]. Pressure values were measured after the interview with a minimum rest of 5 minutes and the person must re-

main seated with their arm resting on the table. Three consecutive measurements were taken with a one-minute interval between one and the other and the systolic and diastolic blood pressure values of the 3 measurements were recorded taking the average into account. Patients were considered hypertensive if systolic BP was ≥ 140 mmHg and diastolic BP was ≥ 90 mmHg [13]. Hypertension was defined according to the diagnostic and classification criteria established by the European Society of Arterial Hypertension, which establishes in adults as Ideal (systolic blood pressure < 120 mmHg and diastolic blood pressure < 85 mmHg); normal (systolic blood pressure < 130 mmHg and diastolic blood pressure < 85 mmHg); borderline (systolic blood pressure 130 mmHg and diastolic blood pressure 85 mmHg or $139 - 89$ mmHg); stage I hypertension (systolic $140 - 159$ mmHg and diastolic $90 - 99$ mmHg); Stage II hypertension (Systolic $160 - 169$ mmHg and diastolic $100 - 109$ mmHg); Stage III hypertension (Systolic ≥ 180 mmHg and diastolic ≥ 110 mmHg); Isolated systolic hypertension (≥ 140 and diastolic < 90 mmHg) [14]. Height and weight were measured with an analogue medical scale containing a non-slip platform, manual adjustment, 600-2100mm altimeter, up to 150kg, 500g graduation with a platform dimension of 270x370, (MEDIKA brand, Portugal). All procedures and standards for reliable results were taken into account considering the manufacturers' descriptions and guidelines. Body mass index was classified according to the WHO classification as normal up to 25, overweight between 25-30 and obese ≥ 30 . After a study was carried out in the cardiology section, patients with high BP were immediately given appropriate treatment.

Statistical Analysis

SPSS version 29 was used to analyze the data. Chi-square test to compare the proportion between hypertensive and non-hypertensive patients. Logistic regression analysis was performed with hypertension as the dependent variable and demographic, behavioural and clinical variables as independent variables. Odds Ratios (OR) and 95% Confidence Intervals (CI) were calculated. All P-values less than 0.05 were considered statistically significant.

Results

The basic characteristics of the participants are shown in Table 1. A total of 158 subjects of both sexes took part in the study, from these 107 (67.7%) were female and 51 (32.7%) males. The age range of the participants varied between 30 and 70 years, with a predominance in the 40s (85.4%) onwards. From the point of view of prevalence, it was found that 67.1% (106/158) of the sample were hypertensive. The mean age and SD among the non-hypertensive subjects was 53.9 ± 13.3 and the hypertensive subjects was 55.8 ± 11.3 . There was no statistical significance in age ($P=0.079$), gender ($P=0.660$), occupation ($P=0.935$), area of residence ($P=0.590$), educational level ($P=0.359$), behavioural characteristics such as physical activity ($P=0.743$) and consumption of alcohol ($p=0.864$) with hypertension. The multivariate logistic regression test showed that patients with hypertension aged between 30 and 40 ($OR=1.19$ CI 95%: 0.3-5.28, $P=0.493$), male ($OR=1.17$ CI 95%: 0.11-13.5, $P=0.660$), employees ($OR=9.94$ CI 95%: 0.48-1.965, $P=0.35$), those living in urban areas ($OR=0.78$ CI 95%:

0.32-1.92%, $P=0.59$), low school graduates ($OR=0.43$ CI 95%: 0.12-1.55%, $P=0.194$), consuming alcohol ($OR=1.11$ CI 95%: 0.33-3.80, $P=0.864$) and practising physical activity ($OR=1.12$ CI 95%: 0.57-2.18, $P=0.742$) had increased risk. In Table 2, we showed the clinical characteristics related to hypertension. Of the 106 patients with hypertension, 81 (76.4%, $P=0.732$) reported having a headache, 25

(23.6%, $P=0.723$) chest pain, 81 (76.4%, $P=0.333$) blurred vision, 63 (59.4%, $P=0.661$) tinnitus and 97 (91.8%, $P=0.189$) nosebleeds. Concerning BMI, the majority (46/106, 43.4%) of hypertensive patients in the sample were obese, followed by those who were overweight (37/106, 34.9%), and (23/106, 21.7%).

Table 1: Demographic and behavioural characteristics related to hypertension among patients from Huambo, Angola.

Independent variables	N (%)	Hypertension			Univariate analysis	
		No (%)	Yes (%)	p-value	OR (95% CI)	p-value
Overall	158 (100)	52 (32.9)	106 (67.1)			
Demographic Characteristic						
Age, year (mean \pm SD)	54.9 \pm 12.0	53.1 \pm 13.4	55.8 \pm 11.3	0.199		
Age distribution						
<30 yo	3 (1.90)	1 (1.90)	2 (1.90)	0.079	0.41 (0.30 - 5.28)	0.493
30 - 40 yo	20 (12.7)	11 (21.2)	9 (8.50)		1.19 (0.11 - 13.5)	0.89
>40 yo	135 (85.4)	40 (76.9)	95 (89.6)		1	
Gender						
Female	107 (67.7)	34 (65.4)	73 (68.9)		1	
Male	51 (32.7)	18 (34.6)	33 (31.1)	0.66	1.17 (0.58 - 2.37)	0.66
Occupation						
Unemployed	104 (65.8)	34 (65.4)	70 (66.0)		1	
Employed	54 (34.2)	18 (34.6)	36 (34.0)	0.935	9.94 (0.48 - 1.95)	0.935
Residence area						
Non-Urbanized	28 (17.7)	8 (15.4)	20 (18.9)	0.59	1	
Urbanized	130 (82.3)	44 (84.6)	86 (81.1)		0.78 (0.32 - 1.92)	0.591
Educational level						
Illiterate	16 (10.1)	4 (10.5)	12 (16.9)	0.359	0.43 (0.12 - 1.55)	0.194
Basic	41 (25.9)	18 (34.6)	23 (21.7)		0.75 (0.21 - 2.69)	0.658
Medium	52 (32.9)	16 (30.8)	36 (34.0)		0.83 (0.23 - 3.03)	0.782
High	52 (47.7)	16 (42.1)	36 (50.7)		1	
Behavioural characteristic						
Alcohol						
No	145 (91.8)	48 (92.3)	97 (91.5)	0.864	1	
Yes	13 (8.20)	4 (7.70)	9 (8.50)		1.11 (0.33 - 3.80)	0.864
Physic activity						
No	70 (44.3)	24 (46.2)	46 (43.4)	0.743	1	
Yes	88 (55.7)	28 (53.8)	60 (56.6)		1.12 (0.57 - 2.18)	0.743

Table 2: Clinical characteristics related to hypertension among patients from Huambo, Angola.

Clinical characteristics	N (%)	Hypertension		
		No (%)	Yes (%)	p-value
Overall	158 (100)	52 (32.9)	106 (67.1)	
Signal and symptoms				
Headache	122 (77.2)	41 (78.8)	81 (76.4)	0.732
Chest pain	36 (22.8)	11 (21.2)	25 (23.6)	0.723
Blurry vision	117 (74.1)	36 (69.2)	81 (76.4)	0.333
Buzz	92 (58.2)	29 (55.8)	63 (59.4)	0.661
Nosebleed	145 (91.8)	48 (92.3)	97 (91.8)	0.864

Asthenia	141 (89.2)	44 (84.6)	97 (91.5)	0.189
Comorbidities				
No	53 (33.5)	17 (32.7)	36 (34.0)	0.874
Yes	105 (66.5)	35 (67.3)	70 (66.0)	
Distribution Comorbidities				
AVC	9 (5.70)	3 (5.80)	6 (5.70)	0.978
Heart failure	48 (30.4)	17 (32.7)	31 (29.2)	0.658
Arterial injury	30 (19.0)	9 (17.3)	21 (19.8)	0.706
Retinal damage	15 (9.50)	97 (91.5)	9 (8.50)	0.539
Aneurysm	1 (0.90)	0 (0.00)	1 (0.90)	0.482
Angina Pectoris	1 (0.60)	0 (0.00)	1 (0.90)	0.482
Body mass index				
Normal	40 (25.3)	17 (32.7)	23 (21.7)	0.011
Overweight	62 (39.2)	25 (48.1)	37 (34.9)	
Obesity	56 (35.4)	10 (19.2)	46 (43.4)	

Discussion

Hypertension has become the main cardiovascular risk factor globally [15]. The data found in the present study showed that hypertension was prevalent in most of the population studied, with about 67% affected by high blood pressure. The prevalence of high blood pressure in the present study was higher than that reported by *Sebastião, et al.* [7-10] in a study addressing the Demographic characteristics and risk factors related to high blood pressure among healthy blood donors from Luanda, the capital city of Angola, where about 7.3% of the population had high blood pressure (>140/90 mmHg) [10].

In an epidemiological cohort study conducted among individuals from Poland, *Zatonska, et al.* (2023) found that several factors played a significant role in increasing the chances of developing hypertension after nine years. These factors included gender, educational level, body weight, alcohol consumption, and embracing a healthy lifestyle [16]. In the present study, patients aged between 30 and 40 (OR=1.19, P=0.493), male (OR=1.17, P=0.660), employees (OR=9.94, P=0.935), patients consuming alcohol (OR=1.11, P=0.864), and practising physical activity (OR=1.12, P=0.742) presented a high likelihood of developing hypertension, while those living in urbanized areas (OR=0.78, P=0.591) and with a low educational level (OR=0.43, P=0.194), had a low likelihood to develop hypertension (Tables 1&2).

In our study, we found that certain factors were associated with a higher likelihood of developing hypertension. Patients who were between the ages of 30 and 40 (OR=1.19, P=0.493), male (OR=1.17, P=0.660), employees (OR=9.94, P=0.935), alcohol consumers (OR=1.11, P=0.864), and those who engaged in physical activity (OR=1.12, P=0.742) had a higher likelihood of developing hypertension. On the other hand, individuals living in urbanized areas and those with a low educational level had a lower likelihood of developing hypertension (Table 1 and Table 2). In previous research, it was discovered that women who are over the age of 40 have a higher likelihood of developing high blood pressure. Interestingly,

our study yielded similar findings, reinforcing this connection [2]. Exercise has been widely recognized as having a remarkable impact on managing high blood pressure and numerous studies have consistently shown this positive effect. However, our study found that physical activity might play a role in contributing to high blood pressure since the prevalence of this condition increased from 54% to 57% among non-hypertensive to hypertensive subjects, respectively [17,18]. The augmented risk of hypertension occurrence among individuals engaged in physical exercise in Angola necessitates comprehensive exploration in future studies. It is imperative to investigate and elucidate the potential underlying reasons behind this association.

Mendy, et al. [19] concludes that the likelihood of having hypertension was significantly higher among those classified as overweight and obese workers compared to their peers [19]. In addition, previous studies carried out in Angola by *Piris, et al.* (2016) using a database from the demographic surveillance system held at the Health Research Centre of Angola (CISA), also concluded that being overweight and obese were significantly associated with hypertension [20]. Our results provide additional evidence that individuals with a normal (22%) body mass index have a reduced likelihood of high blood pressure compared to those who are overweight (35%) or obese (43%), aligning with previous research findings (Table 2). Furthermore, we explored other significant clinical factors, including the occurrence of comorbidities. Our findings revealed that the prevalence of hypertension is twice as high among patients with comorbidities in comparison to those without comorbidities (Table 2).

There are important limitations to this study. The study's statistical power and our ability to draw accurate conclusions are hindered by the small sample size. Additionally, we did not conduct thorough clinical examinations to definitively determine the presence of hypertension. It's important to note that the study's findings may not reflect the overall state of hypertension in Angola, as it was conducted in a specific region. Notwithstanding the limitations inherent in our study, the results unequivocally underscore the

substantial prevalence of cardiovascular disease in Angola, thereby underscoring the urgent necessity for substantial investment in this domain. To this end, we strongly advocate prioritizing research initiatives aimed at identifying high-risk populations prone to developing hypertension. By doing so, we can acquire crucial insights that will inform targeted interventions and ultimately enhance public health outcomes. In summary, hypertension is a significant public health issue in the population of Huambo. It is crucial to implement urgent strategies to prevent the rise in the number of individuals living with high blood pressure.

In this study, we explore the potential demographic factors that may contribute to the rise in hypertension cases in Huambo. It's important to note that the limited number of participants in our study may have affected the statistical power, and the data might not fully represent the actual epidemiological profile of hypertension in the region. Hence, it is crucial to develop effective strategies that focus on enhancing the prevention, diagnosis, and treatment of hypertension. These efforts should extend beyond Huambo and encompass the entire country, to mitigate the rise in cardiovascular diseases among the socially vulnerable population in Angola.

Acknowledgments

The authors thank patients for their participation. We also thank CISA|INIS for the logistical, administrative and scientific support.

Data Availability Statement

The data supporting this study's findings are available on request from the corresponding author.

Conflict Of Interest Statement

The authors declare no conflict of interest.

Author Contribution Statement

Conceptualization: JP and CSS. Investigation: JP, JV, and CSS. Methodology: JP and CSS. Validation: JP, LV, ED, and CSS. Data curation: JP and CSS. Formal analysis: CSS. Data collection: JP. Supervision: LV and ED. Writing - original draft: JP and CSS. Writing - review & editing: JP and CSS. All authors approved the final manuscript for publication.

ORCID

Cruz S Sebastião - 0000-0003-1232-0119.

References

- Shi J, Liu Y, Liu Y, Yi Cheng, Yulu Gu, et al. (2018) Association between ApoE polymorphism and hypertension: A meta-analysis of 28 studies including 5898 cases and 7518 controls. *Gene* 30:675: 197-207.
- Wyatt CM, Chertow GM (2018) Updated guidelines for the diagnosis and management of high blood pressure: implications for clinical practice in nephrology. *Kidney Int* 93(4): 768-770.
- Vetrano DL, Calderón Larrañaga A, Marengoni A, Laura Fratiglioni, Matteo Cesari, et al. (2018) An International Perspective on Chronic Multimorbidity: Approaching the Elephant in the Room. *J Gerontol A Biol Sci Med Sci* 11;73(10): 1350-1356.
- Singer PS (2019) Updates on Hypertension and New Guidelines. *Adv Diatr* 66: 177-187.
- Wu W, Diao J, Yang J, Li Li, Ying Wang, et al. (2021) Impact of Sociodemographic Characteristics, Lifestyle, and Obesity on Coexistence of Diabetes and Hypertension: A Structural Equation Model Analysis amongst Chinese Adults. *Int J Hypertens* 25:2021: 4514871.
- Riley L, Guthold R, Cowan M, Ruth Bonita, Stefan Savin, et al. (2016) The World Health Organization STEPwise Approach to Noncommunicable Disease Risk-Factor Surveillance: Methods, Challenges, and Opportunities. *Am J Public Health* 106(1): 74-8.
- Pires JE, Sebastião YV, Langa AJ, Nery SV (2013) Hypertension in Northern Angola: prevalence, associated factors, awareness, treatment and control. *BMC Public Health* 31: 13:90.
- Paquissi FC, Cuvinje ABP, Cuvinje AB, Paquissi AM (2016) Hypertension among Outpatients at a General Hospital in South Angola: Prevalence, Awareness, Treatment, and Control. *Clin Med Insights Cardiol* 3:10: 111-6.
- Sebastião CS, Sacomboio E, Francisco NM, Joana Morais, Joana Paixão, et al. (2023) Blood pressure pattern among blood donors exposed to SARS-CoV-2 in Luanda, Angola: A retrospective study. *Health Sci Rep* 16;6(8): e1498.
- Sebastião CS, Sacomboio E, Francisco NM, Jocelyne Neto de Vasconcelos, Victor Pimentel, et al. (2023) Demographic characteristics and risk factors related to high blood pressure among healthy blood donors from Luanda, Angola: A retrospective study. *Health Sci Rep* 7;6(6): e1300.
- Gatimu SM, John TW (2020) Socioeconomic inequalities in hypertension in Kenya: a decomposition analysis of 2015 Kenya STEPwise survey on non-communicable diseases risk factors. *Int J Equity Health* 19(1): 213.
- Fonseca Reyes S, Forsyth MacQuarrie AM, García de Alba García JE (2012) Simultaneous blood pressure measurement in both arms in hypertensive and nonhypertensive adult patients. *Blood Press Monit* 17(4): 149-154.
- Vischer AS, Burkard T (2017) Principles of Blood Pressure Measurement – Current Techniques, Office vs Ambulatory Blood Pressure Measurement *Adv Exp Med Biol* 956: 85-96.
- Shi J, Liu Y, Liu Y, Yong Li, Shuang Qiu, et al. (2018) Association between ApoE polymorphism and hypertension: A meta-analysis of 28 studies including 5898 cases and 7518 controls. *Gene* 675: 197-207.
- Parati G, Lackland DT, Campbell NRC, Mayowa Ojo Owolabi, Charlotte Bavuma, et al. (2022) How to Improve Awareness, Treatment, and Control of Hypertension in Africa, and How to Reduce Its Consequences: A Call to Action from the World Hypertension League. *Hypertension* 79(9): 1949-1961.
- Zatońska K, Basiak Rasała A, Połtyn Zaradna K, Dagmara Gaweł Dąbrowska, Maria Wołyniec, et al. (2023) Sociodemographic and behavioral factors associated with controlled hypertension after 9 years of observation of a PURE Poland cohort study. *Front Public Health* 11: 1167515.
- Oliveira GF, Marin TC, Forjaz CLM, Brito LC (2021) Exercise Training and Endothelial Function in Hypertension: Effects of Aerobic and Resistance Training. *Arq Bras Cardiol* 116(5): 948-949.
- Moussouni A, Sidi yakhlef A, Hamdaoui H, Aouar A, Belkhatir D, et al. (2022) Prevalence and risk factors of prehypertension and hypertension in Algeria. *BMC Public Health* 22(1): 1571.
- Mendy VL, Vargas R, Ogungbe O, Zhang L (2020) Hypertension among Mississippi Workers by Sociodemographic Characteristics and Occupation, Behavioral Risk Factor Surveillance System. *Int J Hypertens* 2020: 2401747.
- Pedro JM, Brito M, Barros H (2018) Cardiovascular Risk Assessment in Angolan Adults: A Descriptive Analysis from CardioBengo, a Community-Based Survey. *Int J Hypertens* 2018:2532345.