



Review Article

Copyright© Leonard Sonnenschein

Improving Skin Health: Novel Mechanisms Involving Angiogenesis, Enhanced Topical Delivery, and T-Cell Activation

Leonard Sonnenschein^{1*} and Tony Peticca²

¹The Sonnenschein Institute, 6617 NW 24th Ave, Boca Raton, FL 33496, USA

²Achu Health, 17-235 Industrial Parkway S, Aurora, ON, Canada L4G 3V5

*Corresponding author: Leonard Sonnenschein, The Sonnenschein Institute, 6617 NW 24th Ave, Boca Raton, FL 33496, USA.

To Cite This article: Leonard Sonnenschein* and Tony Peticca, Improving Skin Health: Novel Mechanisms Involving Angiogenesis, Enhanced Topical Delivery, and T-Cell Activation. *Am J Biomed Sci & Res.* 2026 31(6) AJBSR.MS.ID.004092, DOI: [10.34297/AJBSR.2026.31.004092](https://doi.org/10.34297/AJBSR.2026.31.004092)

Received: 📅 July 30, 2026 Published: 📅 August 05, 2026

Abstract

Mild cream- and lotion-based dermal pretreatment supports barrier recovery, hydration, and physiologic conditions conducive to dermal remodeling and angiogenesis, processes linked in the published literature to improved skin texture, barrier function, and appearance. Subsequent topical application of supportive agents maintains a moist healing microenvironment while facilitating delivery of active compounds. Electrocode™, an aqueous mineral-based electricidal solution, has been reported in prior studies to elevate oxygen saturation and stimulate T-cell activation within approximately one hour, coincident with clearance of pathogens and non-viable cells. This review integrates the mechanistic rationale for sequencing cream-based dermal pretreatment with topical Electrocode application to support skin repair and rejuvenation. Independent clinical studies of mild creams and lotions demonstrate that ceramide- and lipid-based formulations accelerate barrier recovery, reduce Transepidermal Water Loss (TEWL), and restore hydration in models of barrier disruption. Electrocode's previously reported effects on oxygenation and rapid T-cell activation may supply a complementary metabolic and immunomodulatory profile relevant to skin barrier integrity and resilience, though this specific application has not been directly studied. The proposed sequence remains a hypothesis-generating framework that requires prospective, controlled clinical evaluation before any efficacy or safety conclusion can be drawn, and no claim of disease treatment or guaranteed cosmetic outcome is made in this review.

Keywords: Skin health, Skin rejuvenation, Angiogenesis, Electrocode, T-cell activation, Barrier recovery, Topical creams, Dermal pretreatment

Introduction

Skin health declines with intrinsic aging, photoaging, chronic inflammation, and impaired barrier function, manifesting as reduced elasticity, uneven texture, diminished vascular support, and increased susceptibility to pathogens. Mild cream- and lotion-based dermal pretreatment supplies supportive lipids and ceramides that accelerate barrier recovery and create favorable conditions for dermal repair pathways, collagen remodeling, and local angiogenesis. Subsequent topical application delivers active compounds efficiently while maintaining moisture and barrier integrity [1].

Electrocode, an aqueous mineral-based electricidal solution developed and studied at The Sonnenschein Institute, has been reported in prior peer-reviewed publications to produce rapid improvements in oxygen saturation and stimulation of T-cell responses in non-dermatological contexts [2-5]. Independent studies of mild topical creams and lotions have shown that lipid- and ceramide-based formulations alone can accelerate barrier repair and improve hydration [6-10]. The present article examines the scientific basis for combining cream-based dermal pretreatment with immediate topical Electrocode application as a potential approach

to support overall skin health, recovery, and long-term dermal integrity. Anecdotal reports have been made observing cut and skin healing occurs rapidly when using health spray. Clinical evaluation is a logical next step for this type of restorative procedure.

Cream-Based Dermal Pretreatment and Support for Skin Health

Mild creams and lotions containing ceramides and physiological lipids support the skin's regenerative capacity by restoring stratum corneum organization, reducing TEWL, and improving hydration. These effects create a favorable microenvironment for keratinocyte and fibroblast activity and can prepare the tissue for improved uptake of subsequently applied topical agents without mechanical disruption.

Angiogenesis as a Core Mechanism Supporting Skin Repair

Healthy skin requires adequate dermal vascularization to deliver oxygen and nutrients to resident cells. During normal wound healing and barrier recovery, angiogenic capillary sprouts organize into a microvascular network, a process mediated by VEGF, FGF, and TGF- β acting on endothelial cell-matrix receptors [1]. Impaired angiogenesis is associated with delayed healing and the visible signs of aged or compromised skin. Cream-based lipid and ceramide support is mechanistically consistent with conditions that favor a pro-angiogenic microenvironment, supporting the vascular requirements of efficient dermal repair; this remains an extrapolation from general wound-angiogenesis and barrier-recovery literature.

Enhanced Topical Delivery and Barrier Support with Cream Approaches

Mild cream and lotion formulations maintain a moist environment, reduce transepidermal water loss, limit microbial colonization during recovery, and deliver regenerative compounds to the layers responsible for barrier restoration and dermal support. Clinical studies of mild creams and lotions have demonstrated measurable barrier healing. A vernix caseosa-based formulation containing ceramides accelerated barrier recovery after tape-stripping, improving TEWL and restoring stratum corneum lipid organization over 16 days [6]. Ceramide-containing creams and lotions in multi-vesicular emulsions produced sustained increases in skin hydration and reductions in dryness lasting at least 24 hours after a single application in dry, eczema-prone skin [7]. Physiological lipid supplementation rebalanced ceramide profiles, strengthened barrier integrity (reduced TEWL after tape-stripping), and decreased irritant sensitivity [8]. A daily ceramide-dominant moisturizing cream and cleanser regimen restored permeability barrier function and improved hydration in moderate eczema [9]. Creams containing ceramides or behenic acid accelerated TEWL recovery in surfactant-damaged skin and enhanced epidermal thickness and density [10]. These findings illustrate that mild topical lipid and ceramide formulations alone can support barrier repair and pro-

vide a mechanistic rationale for their use as dermal pretreatment and subsequent vehicles.

Electrocode: Oxygenation and T-Cell Activation in Support of Skin Health

Electrocode is a hypotonic aqueous solution containing trace minerals reported to exhibit electrical charge properties that selectively affect negatively charged pathogens. In human cohort studies it has produced rapid normalization of subnormal oxygen saturation and body temperature together with improvements in energy, pain, and other self-reported wellness parameters [2].

A dedicated analysis reports that Electrocode administration is associated with an immediate rise in oxygen levels followed by clearance of non-viable cells and pathogens, with resulting products theorized to activate T cells within approximately one hour and initiate antigen and antibody responses [3]. Additional published reports have further characterized Electrocode's antipathogenic activity, immunomodulatory effects, and general wellness support, including a case report describing its adjunct use in an E. coli-associated foodborne illness [4,5].

When applied topically after cream-based pretreatment (as liquid, spray, or cream vehicle), Electrocode may benefit from the optimized barrier environment for local delivery. Its previously reported oxygenation effect may plausibly support the elevated metabolic demands of dermal repair and angiogenesis, while rapid T-cell engagement may plausibly favor controlled regenerative inflammation and skin immune surveillance. These are theoretical extensions of prior findings reported in non-dermatological contexts: no published study has examined Electrocode effects on cutaneous blood flow, collagen synthesis, or validated skin-health outcomes. The barrier-supportive actions of mild creams and lotions documented in independent studies [6-10] further suggest that combining such vehicles with Electrocode could optimize the healing microenvironment.

Proposed Clinical Sequence and Safety Considerations

The following sequence and precautions are presented as an investigational concept intended to inform the design of future controlled studies. They do not constitute a validated treatment protocol, and no claim of disease treatment, permanent rejuvenation, or guaranteed cosmetic outcome is made or implied.

Investigational Protocol Concept

1. Cream-based dermal pretreatment using mild ceramide- or lipid-containing creams or lotions applied evenly to the target skin area (face or body) under clean conditions.
2. Immediate topical application of Electrocode (liquid, spray, or cream formulation), applied evenly to the pretreated area.
3. Optional continued topical-only application of mild bar-

rier-supportive creams or lotions during the recovery phase, consistent with evidence that such formulations accelerate recovery of TEWL and hydration [6-10].

4. Objective, blinded assessment of skin texture, barrier function, elasticity, transepidermal water loss, and safety endpoints over a minimum observation window of 4 weeks, consistent with standard evaluation periods used in dermatologic rejuvenation trials.

Formulation Considerations

Facial and body skin varies substantially in thickness and sensitivity across anatomic sites and must be treated accordingly in any investigational protocol. Mild, non-irritating cream or lotion vehicles designed for cosmetic or barrier-support use are appropriate. Formulations containing harsh exfoliants, strong acids, or high concentrations of irritants are not appropriate and should be excluded from any study protocol.

Safety Considerations

- Restrict use to external, topical application only.
- Exclude participants with broken, bleeding, or actively infected skin, or with active psoriasis, eczema, rosacea flare, or other unstable dermatologic conditions, unless cleared by a licensed dermatologist.
- Exclude skin areas recently treated with chemical peels, laser resurfacing, or other aggressive dermatologic interventions.
- Require patch testing prior to first exposure and monitor for erythema, burning, itching, or hypersensitivity.
- Avoid combination with harsh chemical exfoliants, retinoids, or strong acids on the same day.
- Apply broad-spectrum photoprotection following treatment, consistent with standard aftercare for barrier-support regimens.
- Discontinue use immediately if irritation, rash, or discomfort occurs.

This sequence unites a mechanistically plausible cream-based stimulus for barrier recovery and support of angiogenesis with enhanced topical delivery of an agent that has been reported to supply oxygen support and rapid T-cell engagement; it has not been validated in any clinical trial to date.

Discussion

Cream-based dermal pretreatment supplies an evidence-based approach for barrier recovery, hydration, and conditions that support dermal remodeling and physiologic angiogenesis documented as contributors to improved skin texture and barrier function [1,6-10]. Post-pretreatment topical application supports the healing niche and improved delivery. Incorporation of Electrocode into the

topical step is a logical extension of its previously published effects on oxygenation and T-cell activation, but it remains an untested extrapolation rather than a demonstrated dermatologic effect.

Complementary evidence from studies that used only mild creams and lotions shows that ceramide- and lipid-based formulations reliably accelerate barrier recovery, reduce TEWL, restore hydration, and improve clinical signs of dryness or compromised barrier function [6-10]. These data strengthen the rationale for maintaining a moist, lipid-supported environment and suggest that Electrocode delivered in a compatible mild cream or lotion vehicle could further leverage these barrier-repair mechanisms.

This skin-health application of Electrocode has not been evaluated by the U.S. Food and Drug Administration, Health Canada, or any other regulatory authority for the diagnosis, treatment, cure, mitigation, or prevention of any skin disease or condition. No statement in this review should be interpreted as a clinical, therapeutic, or commercial claim; all proposed mechanisms and outcomes are explicitly hypothesis-generating and require independent, controlled clinical validation before any such claim could be responsibly made.

Prospective randomized controlled trials employing validated skin-assessment scales, transepidermal water-loss measurements, imaging of dermal vasculature, vehicle-only comparator arms, and rigorous safety monitoring are needed.

Conclusion

Cream-based dermal pretreatment followed by topical Electrocode application constitutes a mechanistically coherent approach to supporting skin health. Integration of Electrocode into the topical step leverages its previously reported effects on oxygen levels and rapid T-cell activation to support the angiogenic and regenerative microenvironment favored by barrier recovery. Evidence from independent studies of mild creams and lotions further confirms that barrier-supportive topicals alone can accelerate recovery of TEWL and hydration [6-10], providing additional justification for their inclusion in the sequence. This framework is presented to stimulate rigorous, independent clinical investigation into optimized skin-rejuvenation protocols and should not be interpreted as evidence of therapeutic efficacy.

Conflict of Interest

None.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Ethics Statement

This article is a review of published literature and theoretical synthesis; it does not involve new studies with human participants or animals performed by any of the authors. Therefore, ethical approval was not required. Any future clinical trial testing the se-

quence proposed herein would require institutional review board approval and informed consent.

Acknowledgments

The authors thank Achu Health and The Sonnenschein Institute for institutional support.

References

1. Tonnesen MG, Feng X, Clark RA (2000) Angiogenesis in wound healing. *J Invest Dermatol Symp Proc* 5(1): 40-46.
2. Sonnenschein L, Etyang T, Shah R, Frischer R, Rein G (2021) The Effect of An Aqueous Electricidal Solution on General Well-Being. *Am J Biomed Sci & Res* 13(4): 477-483.
3. Sonnenschein L, Etyang T (2024) T Cell Activation by Natural Mechanisms. *Am J Biomed Sci & Res* 21(6): 661-666.
4. Sonnenschein L, Etyang T, Mwesigye J, Kikonkyo D, Ayella FB, et al. (2024) New Pathways for Anti-Pathogen Responsiveness Using Electrocode. *Am J Biomed Sci & Res* 24(6): 702-716.
5. Sonnenschein L, Etyang T (2025) Unlocking Immunity: ELECTROCODE™ - A Novel Approach. *Am J Biomed Sci & Res* 27(5): 786-790.
6. Boiten WA, Berkers T, Absalah S, van Smeden J, Lavrijsen APM, et al. (2018) Applying a vernix caseosa based formulation accelerates skin barrier repair by modulating lipid biosynthesis. *J Lipid Res* 59(2): 250-260.
7. Danby SG, Andrew PV, Brown K, Chittock J, Kay LJ, et al. (2020) An Investigation of the Skin Barrier Restoring Effects of a Cream and Lotion Containing Ceramides in a Multi-vesicular Emulsion in People with Dry, Eczema-Prone, Skin: The RESTORE Study Phase 1. *Dermatol Ther (Heidelb)* 10(5): 1031-1041.
8. Andrew PV, Williams SF, Brown K, Chittock J, Pinnock A, et al. (2025) Topical supplementation with physiological lipids rebalances the stratum corneum ceramide profile and strengthens skin barrier function in adults predisposed to atopic dermatitis. *Br J Dermatol* 193(4): 729-740.
9. Spada F, Harrison IP, Barnes TM, Greive KA, Daniels D, et al. (2021) A daily regimen of a ceramide-dominant moisturizing cream and cleanser restores the skin permeability barrier in adults with moderate eczema: A randomized trial. *Dermatol Ther* 34(4): e14970.
10. Seo SG, Kwon TG, Ha JW, Oh J, Kim MJ, et al. (2026) Recovery effects of various skin lipid components and their structural derivatives on skin barrier damage. *Int J Cosmet Sci* 48(1): 110-120.